



# ABUNDANCE INDICES FOR NORWEGIAN COASTAL COD NORTH OF 62°N



**Title (English and Norwegian):**

Abundance indices for norwegian coastal cod north of 62°N

Mengdeindekser for kysttorsk nord for 62°N

**Report series:**      **Year - No.:**      **Date:**

Rapport fra havforskningen    2021-6        17.02.2021  
ISSN:1893-4536

**Author(s):**

, Johanna Fall, and Arved Staby (IMR)

Approved by: Research Director(s): Geir Huse Program leader(s): Jan Atle Knutsen

**Distribution:**

Open

**Project No.:**

12352, 15626-06

**Program:**

Kystøkosystemer

**Number of pages:**

105

**Summary (English):**

In connection with the benchmark processes within ICES on Norwegian Coastal Cod north of 62°N during autumn 2020-spring 2021, new abundance index series based on bottom trawl stations and acoustics at the autumn coastal survey were made. The acoustic index covers the period 1995-2019 while the trawl index covers the period 2003-2019. This report describes the methods used and briefly presents the results.

April 30th 2021: Some updates in Appendix.

**Summary (Norwegian):**

I forbindelse med metoderevisjonsarbeidet i ICES på kysttorsk nord for 62°N høsten 2020-våren 2021, ble det utarbeidet nye mengdeindeks-serier basert på faste trålstasjoner og akustikk under kysttoktet om høsten. Akustikkserien dekker perioden 1995-2019, mens trålindeksem dekker perioden 2003-2019. Denne rapporten beskriver metodikken brukt og presenterer kort resultatene i form av tabeller og grafer.

# Content

<b>1</b>	<b>Introduction</b>	5
<b>2</b>	<b>Biology, stock structure, and management of coastal cod</b>	6
<b>3</b>	<b>The autumn coastal survey</b>	7
3.1	The history of the survey and how it has developed over the years	7
3.2	Previous attempts to extract abundance indices from the survey data	9
3.3	Description of the survey data	10
<b>4</b>	<b>Software used</b>	13
<b>5</b>	<b>Acoustic indices</b>	14
5.1	Acoustic indices by length	14
5.2	Acoustic indices by age	15
5.3	Length and weight at age	16
5.4	Uncertainty of abundance indices	16
5.5	Extracting coastal cod from total cod	16
5.6	Acoustic indices - settings in StoX	16
5.7	Resulting time series	18
5.8	Internal consistency in the acoustic series for subarea A	25
5.9	Comparison with acoustic index series calculated with previous methods	27
<b>6</b>	<b>Swept area indices</b>	29
6.1	Swept area indices by length	29
6.2	Swept area indices by age	29
6.3	Length and weight at age	30
6.4	Uncertainty of abundance indices	30
6.5	Extracting coastal cod from total cod	30
6.6	Swept area indices – settings in StoX	31
6.7	Resulting time series	33
6.8	Consistency within the trawl index series for area A	39
<b>7</b>	<b>Comparison of the trawl and acoustic index series</b>	42
<b>8</b>	<b>Conclusions by the data evaluation workshop regarding inclusion of these time series in the assessment of Norwegian Coastal Cod north of 62°N</b>	45
<b>9</b>	<b>References</b>	46
<b>10</b>	<b>Appendix A. Acoustic abundance indices</b>	47
2.1	Total area	47
2.2	Subarea A: North of 67°N	47
2.3	Subarea B: Between 65° and 67°N	47
2.4	Subarea C: Between 62°N and 65°N	47
<b>11</b>	<b>Appendix B Swept area abundance indices</b>	47
3.1	Total area	47
3.2	Subarea A: North of 67°N	47
3.3	Subarea B: Between 65° and 67°N	47
3.4	Subarea C: Between 62°N and 65°N	47
<b>12</b>	<b>Appendix C</b>	93

## 1 - Introduction

New abundance index series for Norwegian coastal cod north of 62° were produced using the software StoX (Johnsen et al. 2019). Indices were calculated by applying trawl swept area and acoustic methods, using data collected during the Norwegian annual coastal survey in autumn NOcoast-Aco-4Q.

Time series of indices for the period 1995 to 2019 in numbers and biomass (2003-2019 for the swept-area index), their coefficient of variation, length- and weight-at-age are tabulated for three subareas and the total area in Appendix A (acoustic indices) and Appendix B (swept-area trawl indices).

## 2 - Biology, stock structure, and management of coastal cod

Coastal cod occurs in fjords and coastal areas along the entire coast of Norway. The management area for this species is divided into two areas south and north of 62 ° N. Genetic studies indicate a genetic cline along the coast from eastern Finnmark to inner Skagerrak (Dahle et al., 2018). Coastal cod north of 62 ° N is somewhat related to Northeast Arctic (NEA) cod, while coastal cod south of 62 ° N showing some similarities to North Sea and Skagerrak cod.

Compared to NEA cod the immature coastal cod has faster individual growth and earlier age of maturation (Berg and Albert, 2003). Since individual growth is reduced after the age of first maturation, the weight at age for old fish is somewhat lower for coastal cod than for Northeast Arctic cod at the same age.

Spawning areas for coastal cod have been mapped by egg sampling and classified according to their relative value (Gyttefelt Torsk MB at [www.fiskeridirektoratet.no](http://www.fiskeridirektoratet.no) ). See also Figures 18-23 in Aglen et al. (2020). Some of these coastal cod spawning areas are close to spawning areas for NEA cod. Probably due to rather small-scale differences in currents and egg buoyancy, the coastal cod eggs and larvae tend to be retained near the spawning areas, while NEA cod eggs and larvae are transported by currents northward into the Barents Sea.

Annual total allowable catches (TAC) were set for coastal cod at 40 kt in the years 1987-2003, 20 kt in 2004, and 21kt in later years. A large proportion of the annual landings of coastal cod is by-catches in the fishery for NEA cod. A rebuilding plan was established 2011 and several technical regulations were introduced to reduce "bycatches" of coastal cod, including gear restrictions and restrictions on vessel size, and closures of spawning areas in the spawning season (Henningsværstraumen and Inner Lofoten).

## 3 - The autumn coastal survey

### 3.1 - The history of the survey and how it has developed over the years

The Institute of Marine Research (IMR) has since 1985 conducted an annual acoustic survey of coastal areas and offshore banks north of 62°N with the objective of obtaining abundance indices of commercially important fish species (Skants, 2019). The annual coverage (in October and November) of coastal areas and fjords, as well as open ocean banks, between Stad (62°N) and Varanger (71.3 °N) has since 1995 included measurements of coastal cod (*Gadus morhua*).

The trawl gear used during the first years was a Campelen 1800 standard shrimp trawl with rock hopper gear and 35 mm mesh size in the cod end. Scanmar sensors provided information about the trawl opening (height in meters), door spread and bottom contact. Since 2003 a Campelen 1800 standard shrimp trawl with rock hopper gear, 20 mm mesh size in the cod end and 80mm (stretched) in the front part is the standard fishing gear, combined with Scanmar trawl and door sensors (Aglen et al., 2005). Additional stations were added in 2017, which was done as it was considered necessary to gather more information on deep water shrimps and redfish (Mehl et al., 2018a). Standard trawl duration is 30 minutes at a speed of 3 knots, with preferred doorspread of 49-52m and trawl opening of 3.5-4.5m. Data were collected with several vessels, which are listed in Table 1.

The survey consists of a stratified grid for acoustic measurements, with fixed bottom trawl stations and additional bottom and pelagic stations on acoustic registrations within each of the strata.

During the surveys hydrographic stations were sampled semi-regularly. CTD-measurements were taken at some of the fixed bottom trawl stations or with a set distance of 30 nautical miles between each station (Staby et al., 2020).

The surveyed area was initially divided into 23 strata, and these were grouped into three subareas: North of 67°N (Area "A"), 65°-67°N (Area "B"), and 62°-65°N (Area "C") (Figure 1). The stratum "Vestfjorden East" was, however, removed from all years, since this stratum had no acoustic coverage and no trawl hauls in most years.

Acoustic transects and bottom trawl hauls are standardized since 2003. In 2017 additional acoustic transects were added to selected strata in order to improve the accuracy of saithe biomass estimates in those strata that contributed a significantly to the total estimate. Figure 2 shows the acoustic transects and trawl hauls made during the coastal survey in 2019.

Trawl catches are sorted and weighed by species according to standard procedures (Mjanger et al., 2020). Length measurements (e.g. total length; from snout to end of the caudal fin) are done for most species, either of all sorted individuals or of a subsample from large catches. Additional information such as age and type from otoliths, sex and gonad maturity stage are collected from cod.





Figure 1. Map showing the 23 strata (lower panel) and the three subareas (upper panel) used for coastal cod index calculations.



Cruise no 2019210 "Johan Hjort". Trawl st.no 704-852. 5.10.-13.11.2019  
Cruise no 2019629 "K. Bonnevie". Trawl st.no 287-406. 2.10.- 9.11.2019

Figure 2. Acoustic transects and trawl hauls made during the cruise in 2019. These are standard transects and trawl hauls that are made during this survey.

### 3.2 - Previous attempts to extract abundance indices from the survey data

As described in chapter 1.2, the autumn coastal survey has a long history and has undergone various changes

over the years. It started out as an acoustic survey primarily targeting coastal cod in the Troms and Finnmark counties in the 1980s. These surveys were conducted by the former "Fiskeriforskning" marine research institute in Tromsø. IMR in Bergen started a survey at approximately the same time of the year, but targeting mainly saithe in the outer coastal areas, partly overlapping the areas of the former survey. A third survey, covering overwintering herring in the fjords was also conducted by IMR. "Fiskeriforskning" was discontinued and the department responsible for resource surveys was included in IMR. From 2003 these surveys were combined and standardized to be a combined acoustic and trawl survey mainly targeting coastal cod and saithe. When the surveys were merged and standardized in 2003, attempts were made to construct an acoustic index for coastal cod based on the original coastal cod survey and the saithe survey, covering also more southern parts of the coast north of 62 ° N. Due to partly different procedures, working protocols, and data format at the two responsible institutes, and since the surveys partly overlapped in space and time, this combination was difficult. The series was started in 1995, but for the years prior to 2003, only some of the acoustic transects, mostly those that had been standardized during the "Fiskeriforskning" survey, were used in index calculations.

### 3.3 - Description of the survey data

Table 1 lists the surveys from 1995-2019, which vessels took part, the number of stations taken etc.

*Table 1. Description of survey data. The number of bottom trawl (BT) stations for swept area and acoustic indices differ because different data filters are applied (see sections 4.1.3 and 4.2.6).*

Year	Vessel	Cruise ID	For swept area index				For acoustic index				
			N BT		N aged	N length	N BT (with cod)	N aged	N length	N miles scrutinized	
			Total	With cod						Total	With cod
1995	Michael Sars	1995111						2515	497	3869	3575
	Johan Hjort	1995211						80	1845	3952	2158
	Volstad	1995810						0	54	0	0
1996	Johan Hjort	1996214						171	2727	3285	1905
	Michael Sars	4-1996						2393	5376	6186	1552
1997	Johan Hjort	1997213						432	3656	3687	2650
	Michael Sars	4-1997						1670	4653	4180	0
1998	G.O.Sars	1998016						493	2237	3030	1808
	Jan Mayen	4-1998						2476	4060	3175	0
1999	Johan Hjort	1999215						399	1083	3813	1653
	Jan Mayen	4-1999						2780	4444	3420	0
2000	Johan Hjort	2000214						414	1202	3766	1816
	Jan Mayen	4-2000						4240	5276	3090	0
2001	Johan Hjort	2001213						267	844	4523	1229
	Jan Mayen	4-2001						3181	3815	2614	0
2002	Johan Hjort	2002214						362	1173	4655	2128
	Jan Mayen	4-2002						2048	2511	3164	0
2003	Johan Hjort	2003211	58	50	1381	2520	72	1580	2808	3695	2012
	Jan Mayen	2003706	78	68	1367	1734	109	1635	2077	4007	3130

2004	Johan Hjort	2004212	50	41	981	1714	67		1159	2010	3502	1989
	Jan Mayen	2004704	81	69	1270	1814	88		1345	1942	3469	2824
2005	Johan Hjort	2005212	42	36	759	1345	51		949	1625	3013	1768
	Jan Mayen	2005704	82	66	695	863	80		689	870	3794	3180
2006	Johan Hjort	2006213	51	42	543	821	48		677	1047	3851	2165
	Håkon Mosby	2006623	0	0	0	0	7		33	51	4479	34
	Jan Mayen	2006705	90	71	865	995	87		886	1021	4013	3260
2007	Johan Hjort	2007212	39	25	179	296	39		327	489	3550	1042
	Håkon Mosby	2007623	0	0	0	0	8		0	103	3778	0
	Jan Mayen	2007703	107	87	834	1040	85		829	1036	4670	3860
2008	Johan Hjort	2008210	86	81	1197	1773	103		1381	2012	6275	5831
	Håkon Mosby	2008623	31	16	233	329	24		308	439	1875	0
	Jan Mayen	2008705	4	4	73	127	10		73	192	214	193
2009	Johan Hjort	2009209	49	41	760	1240	46		818	1306	2860	1696
	Håkon Mosby	2009629	17	6	11	11	5		29	29	1127	180
	Jan Mayen	2009703	66	57	1263	2345	75		1299	2466	2819	2680
	Jan Mayen	2009704	4	4	140	382	14		166	544	524	524
2010	Johan Hjort	2010211	114	90	1957	3847	120		2284	4841	6175	4284
	Jan Mayen	2010704	6	6	149	489	12		168	518	380	380
2011	Johan Hjort	2011214	38	32	536	839	31		561	880	2450	1796
	Helmer Hanssen	2011722	81	71	1246	2075	80		1292	2158	4074	2875
	Helmer Hanssen	2011723	6	6	114	177	18		184	290	240	222
2012	Johan Hjort	2012210	64	55	834	1496	64		974	1701	3760	2552
	Håkon Mosby	2012620	65	51	1129	1878	50		1202	2123	2663	1402
2013	Johan Hjort	2013210	54	51	932	1638	72		1020	1836	3018	1602
	Håkon Mosby	2013623	59	50	1160	2288	65		1249	2451	2339	1227
	Helmer Hanssen	2013851	8	8	124	317	21		124	492	436	432
2014	Helmer Hanssen	2014011	8	8	209	717	21		222	736	455	449
	Johan Hjort	2014213	61	54	928	1601	81		1046	1767	4036	2680
	Håkon Mosby	2014621	74	61	1292	2555	94		1344	2623	2718	1837
2015	Johan Hjort	2015211	64	53	901	1272	65		992	1797	3880	2289
	Håkon Mosby	2015621	74	58	1180	1798	59		1183	2580	2447	1578
	Helmer Hanssen	2015854	8	7	181	566	21		229	613	395	390
2016	Johan Hjort	2016210	70	61	1451	2336	91		1545	2580	4594	4429
	Håkon Mosby	2016620	69	53	976	1463	59		1012	1522	2978	1567
2017	Johan Hjort	2017210	99	89	1616	2822	94		1656	2887	4696	2322
	Kristine Bonnevie	2017620	87	70	917	1854	97		957	1931	3760	1567

2018	Johan Hjort	2018210	110	90	1713	3282	134		1747	3316	4141	1527
	Kristine Bonnevie	2018623	86	66	1301	2365	94		1389	2523	3983	2146
2019	Johan Hjort	2019210	128	100	1516	2059	108		1598	2164	4832	2411
	Kristine Bonnevie	2019629	87	70	1191	2409	117		1196	2499	3345	2371

## 4 - Software used

StoX is a software developed by the Institute of Marine Research for survey analysis and index calculation.

StoX is freely available (<ftp://ftp.imr.no/StoX/Download/>) and is relatively well documented (Johnsen et al., 2019). StoX is currently used for the calculation of bottom trawl indices from the Barents Sea winter survey (Mehl et al., 2018b) and from the Barents Sea ecosystem survey in the autumn (BESS) (Johannesen et al., 2019).

The data was mainly downloaded from: <https://datasetexplorer.hi.no/apps/datasetexplorer/v2/navigation> and the folder "Varanger Stad NOR coastal cruise in autumn». However, for some research vessels, especially in the early part of the period, data was lacking in the relevant folders and had to be retrieved from the original data files and reformatted to the current xml format used by StoX (<https://www.hi.no/hi/forskning/prosjekter/stox>). Steps have been taken to have these data stored in the "Varanger Stad NOR coastal cruise in autumn" folder structure and quality assure for later use.

## 5 - Acoustic indices

A stock abundance index series based on acoustic data from the annual autumn coastal survey ( NOcoast-Aco-4Q ) was calculated using the StoX software (Johnsen et al., 2019). Acoustic data covering the coastline from 62°N to the Russian border are available back to 1995, although the coverage in various parts of this area varied somewhat due to various reasons, see chapter 3.1 and 3.2 for details. For some early years in the series, acoustic data was only available from parts of the survey area. The area was split into 23 strata (see above) and the stock abundance index was calculated for each stratum separately. For various reasons, it was decided to split the total area into three subareas: The coast north of 67°N (A, consisting of 18 strata), between 65° and 67°N (B, consisting of 2 strata), and between 62°N and 65°N (C, consisting of 2 strata) (Fig 1). The spatial coverage during most of the time series is much better in subarea A than in B and C.

To estimate the uncertainty, 500 bootstrap runs were performed, and the indices are the average index from these runs.

### 5.1 - Acoustic indices by length

The conversion of mean nautical area scattering coefficient (NASC,  $m^2 \text{ nmi}^{-2}$ ) to fish density was carried out using a standard procedure, where trawl stations (with a catch of more than 1 cod individual) were assigned to each PSU. As a rule, all stations within a stratum were assigned to the PSUs in the same stratum. However, if less than three trawl stations were carried out in a stratum, stations in neighbouring strata were assigned to the PSUs so that at least three stations were assigned to each PSU.

The combined length distribution ( $d$ ), calculated for each transect (PSU,  $j$ ), is given by

$$d_{l,j} = \sum_{s=1}^n d_{l,s,j}, \quad (\text{eqn 1})$$

where  $d_{l,s,j}$  is fish density (number by 1 nmi tow distance) by station ( $s$ ) and length group ( $l$ , cm), and  $n$  is the total number of stations.

The fish density ( $\rho$ , individuals  $\text{nmi}^{-2}$ ) by length group and transect was calculated using

$$\rho_{j,l} = \text{NASC}_{j,l} / (4\pi\sigma_{bs,l}), \quad (\text{eqn 2})$$

where  $\text{NASC}_{j,l}$  is the mean nautical area scattering coefficient by transect and length group and  $\sigma_{bs,l}$  ( $\text{m}^2$ ) is the acoustic backscattering cross-section for a fish of length  $l$ .

$\text{NASC}_{j,l}$  is given by

$$\text{NASC}_{j,l} = \text{NASC}_j p_{l,j} (\sigma_{bs,l} / (\sum_l \sigma_{bs,l})), \quad (\text{eqn 3})$$

where  $\sigma_{bs,l}$  is the acoustic backscattering cross-section for a fish of length  $l$  multiplied by the proportion ( $p$ ) of a fish of length  $l$  in  $d_{l,j}$ , and  $\text{NASC}_j$  is the mean nautical area scattering coefficient over a given transect.

The acoustic backscattering cross-section for a fish of length  $l$  is calculated using

$$\sigma_{bs,l} = 10^{\left(\frac{TS_l}{10}\right)}, \text{ (eqn 4)}$$

where the target strength, TS (dB re 1m<sup>-2</sup>), for a fish of length l is calculated using

$$TS_l = m \log_{10}(l) + a, \text{ (eqn 5)}$$

where m and a are constants, set at 20 and -68.0 respectively.

The abundance (N, inds) of cod by length group l and stratum k is given by

$$N_{k,l} = \rho_{k,l} A_k, \text{ (eqn 6)}$$

where A (nmi<sup>2</sup>) is the stratum area, and the mean density of cod by l and k is given by

$$\rho_{k,l} = (1/n_k) \sum_{j=1}^{n_k} (w_{k,j} \rho_{k,j,l}), \text{ (eqn 7)}$$

where  $\rho_{k,l} = (1/n_k) \sum_{j=1}^{n_k} (w_{k,j} \rho_{k,j,l})$ , is the transect weight,  $n_k$  is the total number of sample transects and  $L_{k,j}$  and  $\bar{L}_k$  are the distance of each transect by stratum and the mean transect distance over each stratum respectively.

## 5.2 - Acoustic indices by age

Only a subsample of the length-measured individuals (j) is aged. A two-stage conversion process is used to convert the abundance of fish by length group to abundance of fish by age group.

First, the abundance ( $N_{k,l}$ ) by l and k is distributed the length-measured individuals to generate so-called super-individuals, each representing an abundance estimated as:

$$N_{k,j,s,l} = N_{k,l} w_{k,j,s,l}, \text{ (eqn 8)}$$

where

$$w_{k,j,s,l} = \rho_{k,s,l} / \left( \sum_{s=1}^n \rho_{k,s,l} \right) \times 1/m_{k,s,l}, \text{ (eqn 9)}$$

and  $m$  is the number of length-measured individuals.

Second, in instances where a super-individual is not aged, the missing age is filled in by a random data imputation. The imputation of missing age is principally carried out at the station level, randomly selecting the value from aged super-individuals within the same length group. If no aged super-individual is available at station level, the imputation is attempted at strata level, or lastly on survey level. In instances, where no age information is available at any level for a specific length group, the abundance estimate is presented with unknown age. As the imputation of missing age values in both examples also imputes associated biological parameters, abundance can be estimated for any combination of classifications assigned to the super-individuals e.g. sex, maturity, age etc. In our case the otolith type was used to classify the super-individuals, see below.

### 5.3 - Length and weight at age

Length and weight at age was calculated using the weighting factors defined in eqn 8 (the “super -individuals”).

### 5.4 - Uncertainty of abundance indices

Uncertainty was estimated as the coefficient of variation (ratio of standard deviation to the mean, CV). StoX calculates CV using bootstrap runs by stratum, treating each trawl station as the primary sampling unit. Here we used 500 bootstrap runs.

### 5.5 - Extracting coastal cod from total cod

Since the discrimination of coastal cod and other cod caught at the coastal survey is based on otolith types (see above), this poses a special challenge to producing abundance index series with uncertainty for coastal cod. Running a StoX project on the acoustical and biological data to produce an acoustic index series will primarily produce indices for all cod present in these data sources. However, when running the bootstrap process in StoX, it is possible to group the superindividuals by several categories, for instance age and otolith type. There is no facility inside StoX to present those “two-dimensional” bootstrap data but using an R-script manipulating the bootstrap files generated by StoX it is possible to extract relevant data. Thus, this was done after the whole time series was made by ordinary StoX-runs, by selecting only those entries in the bootstrap data that contained superindividuals with otolith types “1” and “2”. All tables and figures in the appendices were produced with this R-script. The R-script itself is documented in appendix C. Since the growth pattern can only be distinguished with certainty in otoliths from two-year old and older fish (although an otolith type is in some cases noted also for younger fish), the indices of age 0 and age 1 were excluded from the index series suggested for use in stock assessment.

### 5.6 - Acoustic indices - settings in StoX

The processes included and the settings of parameters when running StoX for acoustic indices are given in Tables 2-5:

Table 2. Baseline processes:

Process	Parameters	Values
<b>ReadProcessData</b>		
<b>ReadAcousticXML</b>	FileName1, FileName2, ...	Relevant data files
<b>FilterAcoustic</b>	AcousticData	ReadAcousticXML
	DistanceExpr	N/A
	FreqExpr	N/A
	NASCExpr	acocat == 31
<b>NASC</b>	LayerType	WaterColumn
<b>ReadBioticXML</b>	FileName1, FileName2, ...	Relevant file names
<b>FilterBiotic</b>	FishStationExpr	fs.getLengthSampleCount('TORSK')>1
	CatchExpr	species == '164712'
	SampleExpr	N/A

	IndExpr	N/A
<b>StationLengthDist</b>	LengthDistType	NormLengthDist
<b>RegroupLengthDist</b>	LengthInterval	1.0
<b>Catchability</b>	CatchabilityMethod	LengthDependentSweepWidth
	LengthDist	RegroupLengthDist
	ParLengthDependentSweepWidth	SpecCat=;Alpha=5.91;Beta=0.43;LMin=15.0;LMax=62.0
<b>RelLengthDist</b>	LengthDist	Catchability
<b>DefineStrata</b>	UseProcessData	" True"
<b>StratumArea</b>	AreaMethod	Accurate
<b>DefineAcousticTransect</b>	DefinitionMethod	UseProcessData
<b>MeanNASC</b>	NASC	NASC
	SampleUnitType	PSU
<b>BioStationAssignment</b>	BioticData	FilterBiotic
	AssignmentMethod	Stratum (first time, then UseProcessData)
	EstLayers	1~PELBOT
<b>BioStationWeighting</b>	WeightingMethod	SumWeightCount
<b>TotalLengthDist</b>	LengthDist	RelLengthDist
<b>AcousticDensity</b>	LengthDist	TotalLengthDist
	NASC	MeanNASC
	m	20
	a	-68
<b>MeanDensity_Stratum</b>	Density	AcousticDensity
	SampleUnitType	Stratum
<b>SumDensity_Stratum</b>	Density	MeanDensity_Stratum
<b>Abundance</b>	Density	SumDensity_Stratum
	PolygonArea	StratumArea
<b>IndividualDataStations</b>	Abundance	Abundance
<b>IndividualData</b>	IndividualDataStations	IndividualDataStations
<b>SuperIndAbundance</b>	Abundance	Abundance
	IndividualData	IndividualData
	AbundWeightMethod	StationDensity
	LengthDist	RegroupLengthDist

Table 3. Baseline report processes:

Process	Parameters	Values
<b>FillMissingData</b>	Superindividuals	SuperIndAbundance
	FillVariables	ImputeByAge

	Seed	1
	FillWeight	Mean
<b>EstimateByPopulationCategory</b>	Superindividuals	FillMissingData
	Lengthinterval	5.0
	Scale	1000
	Dim1	otolithtype
	Dim2	age
	Dim3	SpecCat

Table 4. R processes:

Process	Parameters	Values
<b>runBootstrap</b>	bootstrapMethod	AcousticTrawl
	acousticMethod	PSU~Stratum
	bioticMethod	PSU~Stratum
	startProcess	TotalLengtDist
	endProcess	SuperIndAbundance
	nboot	500
	seed	1
	cores	4
<b>imputeByAge</b>	seed	1
	cores	4
<b>SaveProjectData</b>		“ Enabled”

Table 5 R report processes:

Process	Parameters	Values
<b>getReports</b>	out	all
	options	grp1="age", grp2="otolithtype"
<b>getPlots</b>	out	all
	options	grp1="age", grp2="otolithtype"

## 5.7 - Resulting time series

The annual abundance indices and biomass indices by age groups and for age group 2+, their coefficient of variation, and mean length and weight by age groups are shown in Appendix A for the total area and for the subareas A, B and C.

The abundance indices for age 2+ are depicted in Figures 3 to 6. The series for the total area (Fig 3) is characterized by high indices at the beginning of the time series, but rapidly decreasing from 1997 to a level of 10-20 million, without any clear trends. In general, the uncertainties are larger during the first part of the time series compared to more recent years.




Figure 3. Acoustic index series for coastal cod age 2+ in the total area. Error bars represent +/- two standard deviations.

The series for subarea A (the northern part of the survey area, Figure 4) resembles that for the total area, because this area contains most of the coastal cod.




Figure 4. Acoustic index series for coastal cod age 2+ in subarea A (north of 67 ° N). Error bars represent +/- two standard deviations.

The indices for subareas B (Figure 5) and C (Figure 6) are lower than for subarea A. The uncertainties are also more variable from year to year.




Figure 5. Acoustic index series for coastal cod age 2+ in subarea B (between 65 ° N and 67 ° N). Error bars represent +/- two standard deviations.




Figure 6. Acoustic index series for coastal cod age 2+ in subarea C (between 62 °N and 65 °N). Error bars represent +/- two standard deviations.

To assess whether the indices can describe the stock dynamics over time, plots on how year classes (cohorts) could be traced from year to year were constructed (Figure 7-10). The progression of year classes through the stock is reasonably well described for the total area and for subarea A (Figures 7-8). A year effect is visible for instance in 1998, when all age groups where recorded lower than expected. In other years, single age groups, in particular among older fish, show unexpected patterns. This seems to be a problem for age groups above 10 years. As expected, the plot for Subarea A (Figure 8) resembles that for the total area, while those for Subarea B (Figure 9) and Subarea C (Figure 10) show a much less consistent picture, where strong year effects are visible .




Figure 7. Abundance (log scale) over time for the year classes 1995 to 2014 for age 1 and older in the acoustic index series for the total area. The age is shown for each data point.




Figure 8. Abundance (log scale) over time for the year classes 1995 to 2014 for age 1 and older in the acoustic index series for Subarea A. The age is shown for each data point.





Figure 9. Abundance (log scale) over time for the year classes 1995 to 2014 for age 1 and older in the acoustic index series for Subarea B. The age is shown for each data point.






Figure 11. Consistency plots for the acoustic index for area A. Age groups 1-6.




Figure 12. Consistency plots for the acoustic index for area A. Age groups 7-12.

## 5.9 - Comparison with acoustic index series calculated with previous methods

An acoustic index has been calculated for this stock since 1995 (see chapter 1.3). Because this approach used a somewhat more detailed strata system and a method based on the SAS software platform, it is difficult to compare that series with the new acoustic index developed in StoX. Nonetheless, a comparison made for the sum over ages 2+ (Figure 13) shows that there are large differences between the series before 2002. For the period after 2002 the indices are much more similar, with the new series estimates being somewhat lower in most years. A part of the large discrepancies found in the early years of the series can be explained by poor data quality of the acoustic data. When the series was rerun in StoX and the acoustic data checked in detail, some few enormous NASC values were detected, indicating that parts of the bottom signal had been integrated. In small strata containing few transects and few values, such erroneous values may have a big impact on the total index. For instance, in 2000, taking out a NASC value of 4718 from a small stratum (Sørøya Indre) made the index in that stratum change from 46 000 tonnes to 1400 tonnes. In 1996, 1999 and in 2000 six extremely high values were found and removed from the transects, which had profound effects on the index values from these years. It is unknown whether these erroneous values were in fact included in the old acoustic series or not. In the remaining years before 2002, no extreme values could be found and also in these years the new index is substantially lower than the old index, indicating that there are other reasons for the differences.

Despite this we argue that the new indices should be accepted on the grounds that they are developed in one go, using a more quality assured software with identical settings from year to year, and with a more thorough quality assurance of the acoustic data.




Figure 13. Comparison of the old and new acoustic abundance index series for sum over ages 2+ for the total area.

## 6 - Swept area indices

A stock abundance index series based on bottom trawl hauls at the annual autumn coastal survey ( NOcoast-Aco-4Q ) was calculated using the StoX software. Trawl data covering the coastline from 62°N to the Russian border were available back to 1995, although the coverage in various parts of this area varied somewhat due to ship availability, weather conditions etc. However, the survey was designed as an acoustic survey before 2003, and trawls were done on acoustic registrations in order to supply biological data for the acoustic observations. Consequently, we did use trawl data prior to 2003 when calculating the swept-area indices for coastal cod. The area is split into 22 strata (Figure 1) and the stock abundance index is calculated for each stratum separately. It was decided to split the total area into three subareas: The coast north of 67°N (A, consisting of 18 strata), between 65° and 67°N (B, consisting of 2 strata), and between 62°N and 65°N (C, consisting of 2 strata). The coverage during most of the time series is much better in subarea A than in B and C.

The coastal cod was split from the NEA cod using the same methods as described for the acoustic index, see above.

To estimate the uncertainty of the estimate, 500 bootstrap runs were performed, and the indices are the average index from these runs.

### 6.1 - Swept area indices by length

The following description is taken from Johannessen et al. (2019):

The swept area density ( $\rho$ , individuals per square nautical mile, inds nmi<sup>-2</sup>) by stratum (k), station (s) and length group l (1 cm), is given by

$$\rho_{k,s,l} = f_{k,s,l} / sw_l, \text{ (eqn 10)}$$

where  $f_{k,s,l}$  is the number of individuals standardized over a towing distance of 1 nmi by k, s and l, and  $sw_l$  is the adjusted swept width in nmi's by length group calculated using

$$sw_l = EW_l / 1852, \text{ (eqn 11)}$$

where  $EW_l$  is the length dependent effective swept width. The length dependency of swept width is taken from (Dickson, 1993)

The abundance ( $N$ , inds) by l and k is calculated using

$$N_{k,l} = \rho_{k,l} A_{k,l}, \text{ (eqn 12)}$$

where  $A$  is stratum area (nmi<sup>2</sup>), and  $\rho_{k,l}$  is the average swept area density by l and k, given by

$$\rho_{k,l} = [1/n] \sum_{s=1}^n \rho_{k,s,l}, \text{ (eqn 13)}$$

where n is number of stations.

### 6.2 - Swept area indices by age

The sampling protocol for the survey is to sample one individual from each 5 cm length group at each trawl

station for aging and individual weights. A two-stage conversion process is used to convert the abundance of fish by length group to abundance of fish by age group.

Firstly, the abundance ( $N_{k,l}$ ) by length group l (5 cm) and stratum k is distributed by the length-measured individuals (j) to generate so-called “Super-individuals” (super-individuals represent fractions of a total, our use corresponds to a probability based design where  $w_{k,j,s,l}$  is the inverse of the inclusion probability for a single fish sample), each representing an abundance estimated as:

$$N_{k,j,s,l} = N_{k,l} w_{k,j,s,l}, \text{ (eqn 14)}$$

where

$$w_{k,j,s,l} = \frac{\rho_{k,s,l}}{\left( \sum_{s=1}^n \rho_{k,s,l} \right)} \times \frac{1}{m_{k,s,l}}, \text{ (eqn 15)}$$

and  $m$  is the number of length-measured individuals

Secondly, in instances where a super-individual is not aged, the missing age is filled in by a random data imputation. The imputation of missing age is principally carried out at the station level, randomly selecting the value from aged super-individuals within the same length group. If no aged super-individual is available at the station level, the imputation is attempted at strata level, or lastly on survey level. In instances where no age information is available at any level for a specific length group, the abundance estimate is presented with unknown age (Johnsen et al., 2019).

### 6.3 - Length and weight at age

Length and weight at age was calculated using the weighting factors defined in eqn 15 (the “super-individuals”).

### 6.4 - Uncertainty of abundance indices

Uncertainty was estimated as the coefficient of variation (ratio of standard deviation to the mean, CV). StoX calculates CV using bootstrap runs by stratum, treating each trawl station as the primary sampling unit. Here we used 500 bootstrap runs.

### 6.5 - Extracting coastal cod from total cod

Since the discrimination of coastal cod and other cod caught at the coastal survey is based on otolith types (see above) this poses a special challenge to producing abundance index series with uncertainty for coastal cod. Running a StoX project on the biological data to produce a swept-area index series will primarily produce indices for all cod present in this data source. However, when running the bootstrap process in StoX, it is possible to group the superindividuals by several categories, for instance age and otolith type. There is no facility inside StoX to present those “two-dimensional” bootstrap data but using an R-script manipulating the bootstrap files generated by StoX it is possible to extract relevant data. Thus, this was done after the whole time series were made by ordinary StoX-runs, by selecting only those entries in the bootstrap data that contained superindividuals with otolith types “1” and “2”. All tables and figures in the appendices to this document were produced by this R-script. The R-script itself is documented in appendix “C”.

## 6.6 - Swept area indices – settings in StoX

The processes included and the settings of parameters when running StoX for swept area indices are given in tables 6-9:

Table 6. Baseline processes:

Process	Parameters	Values
<b>ReadProcessData</b>		
<b>ReadBioticXML</b>	FileName1, FileName2, ...	Relevant file names
<b>FilterBiotic</b>	FishStationExpr*	gear =~['3270','3271'] and gearcondition < 3 and tawlquality =~['1','3'] and fishstationtype !=['2'] and
	CatchExpr	species == '164712'
	SampleExpr	N/A
	IndExpr	N/A
<b>DefineSweptAreaPSU</b>	Method	Station
<b>StationLengthDist</b>	LengthDistType	NormLengthDist
<b>RegroupLengthDist</b>	LengthInverval	5.0
<b>Catchability</b>	CatchabilityMethod	LengthDependentSweepWidth
	LengthDist	RegroupLengthDist
	ParLengthDependentSweepWidth	SpecCat=;Alpha=5.91;Beta=0.43;LMin=15.0;LMax=62.0
<b>RelLengthDist</b>	LengthDist	Catchability
<b>DefineStrata</b>	UseProcessData	" True"
<b>StratumArea</b>	AreaMethod	Accurate
<b>TotalLengthDist</b>	LengthDist	RegroupLengthDist
<b>SweptAreaDensity</b>	SweptAreaMethod	LengthDependent
	BioticData	FilterBiotic
	LengthDist	TotalLengthDist
	DistanceMethod	FullDistance
	SweepwidthMethod	Predetermined
<b>MeanDensity_Stratum</b>	Density	SweptAreaDensity
	SampleUnitType	Stratum
	PolygonArea	StratumArea
<b>AbundanceByLength</b>	Density	MeanDensity_Stratum
<b>IndividualDataStations</b>	Abundance	AbundanceByLength
<b>IndividualData</b>	IndividualDataStations	IndividualDataStations
<b>SuperIndAbundance</b>	Abundance	AbundanceByLength
	IndividualData	IndividualData

	AbundWeightMethod	StationDensity
	LengthDist	RegroupLengthDist

\* In the period 2017-2019 this filter was changed to allow for inclusion of stations coded with StationType = C and trawlQuality = 2

Table 7. Baseline report processes:

Process	Parameters	Values
<b>FillMissingData</b>	Superindividuals	SuperIndAbundance
	FillVariables	ImputeByAge
	Seed	1
	FillWeight	Mean
<b>EstimateByPopulationCategory</b>	Superindividuals	FillMissingData
	Lengthinterval	5.0
	Scale	1000
	Dim1	otolithtype
	Dim2	age
	Dim3	SpecCat

Table 8. R processes:

Process	Parameters	Values
<b>runBootstrap</b>	bootstrapMethod	AcousticTrawl
	acousticMethod	PSU~Stratum
	bioticMethod	PSU~Stratum
	startProcess	TotalLengtDist
	endProcess	SuperIndAbundance
	nboot	500
	seed	1
	cores	4
<b>imputeByAge</b>	seed	1
	cores	4
<b>SaveRImage</b>		“ Enabled”

Table 9. R report processes:

Process	Parameters	Values
<b>getReports</b>	out	all
	options	grp1="age", grp2="otolithtype"
<b>getPlots</b>	out	all

	options	grp1="age", grp2="otolithtype"
--	---------	--------------------------------

## 6.7 - Resulting time series

Index series for the total area (Figure 14) and for the three subareas A (Figure 15), B (Figure 16) and C (Figure 17) are shown below. The abundance indices for the total area is rather flat but one year, 1997, stands out from the rest having a three times as high index and much wider confidence limits than the rest of the years in the series.

The amount of coastal cod in subarea A (Figure 15) is much higher than in the more southern subareas B and C, and the index series in subarea A therefore resembles the total index to a high degree. While the index for subarea A (and therefore also the total area) shows peaks in 2003 and in 2014-2015, the series for subareas B and C are without conspicuous trends. The relative uncertainty is much higher for the two southern subareas than for the northern (subarea A). The uncertainty in the last four years is smaller than for the earlier part of the index series, for all subareas.




Figure 14. Trawl index series for coastal cod age 2+ in the total area. Error bars represent +/- two standard deviations.




Figure 15. Trawl index series for coastal cod age 2+ in subarea A (north of 67 ° N). Error bars represent +/- two standard deviations.




Figure 16. Trawl index series for coastal cod age 2+ in subarea B, (between 65 ° N and 67 ° N). Error bars represent +/- two standard deviations.




Figure 17. Trawl index series for coastal cod age 2+ in subarea C, (between 62 ° N and 65 ° N). Error bars represent +/- two standard deviations.

Consistency among cohorts are illustrated on Figures 18-21. The cohorts can be followed in subarea A (and in the total area) without conspicuous year effects, except in 2005 and 2007. For the two subareas B (Figure 20) and C (Figure 21) it is not possible to follow the year classes except for short periods, indicating that the indices for these subareas do not reflect the total abundance of coastal cod during the period 2002 to 2019.





Figure 18. Abundance (log scale) over time for the year classes 2002 to 2014 for age 1 and older in the trawl index series for the total area. The age is shown for each data point.






Figure 20. Abundance (log scale) over time for the year classes 1995 to 2014 for age 1 and older in the trawl index series for Subarea B.

B. The age is shown for each data point.




Figure 21. Abundance (log scale) over time for the year classes 2002 to 2014 for age 1 and older in the trawl index series for Subarea C. The age is shown for each data point.

## 6.8 - Consistency within the trawl index series for area A

The internal consistency plots (number at age n in year n plotted versus number at age n+1 in year y+1) for age groups 1-6 are shown in Figure 22 and for age groups 7-12 in Figure 23. In most cases the fit is poor.




Figure 22. Consistency plots of the trawl index for area A. Age groups 1-6.




Figure 23. Consistency plots of the trawl index for area A. Age groups 7-11.

## 7 - Comparison of the trawl and acoustic index series

The acoustic index series and the trawl index series give a partly independent view of the stock situation over time. They are not totally independent, since the length information used to translate the acoustic backscatter into fish abundance comes by and large from the same trawl hauls that are used for calculation of swept area indices, and the age information used to break the acoustic index down to age groups partly comes from the same trawl hauls that are used to calculate the swept area indices. However, the total backscatter, mainly determining the acoustic index, is totally independent of the catch rates in the trawl hauls, so in this respect the two series give independent information about the amount of fish. There are numerous reasons that these indices differ. Trawling on the bottom is only possible where the bottom is trawlable, that is soft and smooth and not too steep. In many areas of the coast it is not possible to trawl, and consequently the trawl hauls may not be representative of areas with hard and/or steep bottom. On the other hand, even though the acoustic method will cover all navigable waters, the acoustic backscatter signal is difficult to interpret where the bottom is steep, and in all areas the dead zone near the bottom will not be covered. Also, the allocation of acoustic backscattering by species can be challenging, and in particular in areas where it is not possible to trawl.

In Figure 24 the new acoustic index series and the swept-area series are compared, and also the landings statistics are included on the figure for comparison.




Figure 24. New acoustic series (2+), trawl index series (2+) and landings (taken from table 2.1a in ICES (2020)).

The acoustic series, going back to 1995, shows a decrease in the last part of the 1990s with a simultaneous decrease in catches during that period, from a record high catch in 1995 to a level at about half of that total during the next decades. The acoustic index and the trawl index fluctuate without clear trends after 2003, in some years the acoustic index is higher than the trawl index and vice versa. These index series are compared on age-group basis in figure 25 and 26. The consistency is quite good for many of the age groups, with  $r^2$  in the

range 0.2-0.6. However, for some age groups (mainly 3-6) the fit is poorer, with  $r^2$  in the range 0.0-0.1.




Figure 25. Comparison of acoustic index and trawl index for area A in the period 2003 to 2009, age groups 1-6.




Figure 26. Comparison of acoustic index and trawl index for area A in the period 2003 to 2009, age groups 7-12.

## **8 - Conclusions by the data evaluation workshop regarding inclusion of these time series in the assessment of Norwegian Coastal Cod north of 62°N**

Based on an evaluation of what is presented in this document concerning data quality, survey coverage, and year-to-year consistency, the data evaluation workshop held in December 2020 concluded that

1. The acoustic abundance index series from 1995 to 2019, for age groups 2-10+, may be used as input data in analytical assessment models for coastal cod in subarea A. The corresponding estimates of length- and weight-at-age may be used as estimates of length- and weight-at-age in the stock.
2. For subareas B and C, the acoustic indices for biomass of age 2+ may be used in biomass models or to assess changes in stock abundance from year to year, using methods for data-limited stocks.
3. The trawl index series from 2003 to 2019, for age groups 2-10, may be used as input data in analytical assessment models for coastal cod in subarea A. The corresponding estimates of length- and weight-at-age may be used as estimates of length- and weight-at-age in the stock.
4. For subareas B and C, the trawl indices of biomass of age 2+ may be used in biomass models or to assess changes in stock abundance from year to year, using methods for data-poor stocks.

## 9 - References

- Aglen, A., Berg, E., Mehl, S., and Sunnanå, K. 2005. Acoustic abundance of saithe, coastal cod and juvenile herring Finnmark – Møre Autumn 2005. Toktrapport/Havforskningsinstituttet, 2005/23.
- Aglen, A., Nedreaas, K., Knutsen, J. A., and Huse, G. 2020. Kysttorsk nord for 62 grader nord. Vurdering av status og forslag til forvaltingstiltak og ny gjenoppbyggingsplan. Fisken og Havet, 2020/2: 1-64.
- Berg, E., and Albert, O. T. 2003. Cod in fjords and coastal waters of North Norway: distribution and variation in length and maturity at age. ICES Journal of Marine Science, 60: 787-797.
- Dahle, G., Quintela, M., Johansen, T., Westgaard, J. I., Besnier, F., Aglen, A., Jorstad, K. E., et al. 2018. Analysis of coastal cod (*Gadus morhua* L.) sampled on spawning sites reveals a genetic gradient throughout Norway's coastline. BMC Genet, 19: 42.
- Dickson, W. 1993. Estimation of the capture efficiency of trawl gear. 1:Development of a theoretical model. Fisheries Research, 16: 239-253.
- ICES 2020. Arctic Fisheries Working Group (AFWG). ICES Scientific Reports., 2:52. : 577
- Johannesen, E., Johnsen, E., Johansen, G. O., and Korsbrekke, K. 2019. StoX applied to cod and haddock data from the Barents Sea NOR-RUS ecosystem cruise in autumn. Swept area abundance, length and weight at age 2004-2017. Fisken og Havet, 2019-6: 1-40.
- Johnsen, E., Totland, A., Skålevik, Å., Holmin, A. J., Dingsør, G. E., Fuglebakk, E., Handegard, N. O., et al. 2019. StoX: An open source software for marine survey analyses. Methods in Ecology and Evolution, 10: 1523-1528.
- Mehl, S., Aglen, A., Berg, E., Dingsor, G. E., and Korsbrekke, K. 2018a. Akustisk mengdemåling av sei, kysttorsk og hhyse Finnmark-Møre hausten 2017. Toktrapport/Havforskningsinstituttet, Nr. 2–2018.
- Mehl, S., Aglen, A., Johnsen, E., and Skålevik, Å. 2018b. Estimation of acoustic indices with CVs for cod and haddock in the Barents Sea winter survey 1994 – 2017 applying the Sea2Data StoX software. Fisken og Havet, 5-2018.
- Mjanger, H., Svendsen, B. V., Senneset, H., Fuglebakk, E., Skage, M. L., Diaz, J., Johansen, G. O., et al. 2020. Handbook for sampling fish, crustaceans and other invertebrates. pp. 1-157. Institute of Marine Research, Bergen.
- Skants, K. D. 2019. Species composition, distribution and ecology of the demersal fish community along the Norwegian coast north of Stad under varying environmental conditions. In Department of Biological Sciences - Marine Biology. University of Bergen, Norway.
- Staby, A., Mehl, S., Berg, E., Aglen, A., and Korsbrekke, K. 2020. Acoustic abundance of saithe and coastal cod Finnmark – Møre Autumn 2018. Toktrapport/Havforskningsinstituttet, Nr. 5–2020.

## 10 - Appendix A. Acoustic abundance indices

### 2.1 - Total area

Table A.1.1. Abundance indices (millions)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17
1995	0.031	26.961	11.015	7.254	7.207	7.023	4.618	1.561	0.678	0.340	0.042		0.134					
1996	22.040	17.763	10.743	12.628	6.669	7.434	3.386	1.269	0.213	0.034	0.123							
1997	0.018	17.724	17.907	20.326	9.288	5.243	2.652	0.919	0.393	0.059	0.014	0.019						
1998	1.269	8.713	10.675	10.731	9.626	4.238	1.806	0.951	0.141	0.123	0.037		0.009		0.026	0.026		
1999		2.562	3.990	4.112	3.283	2.794	0.941	0.214	0.030	0.067	0.015		0.005					
2000	1.979	5.264	8.468	7.426	4.935	4.320	3.106	0.712	0.307	0.087	0.029	0.026		0.010				
2001	0.207	2.725	4.847	4.734	4.343	2.516	1.637	1.018	0.219	0.031	0.036	0.029	0.009	0.018				
2002	0.418	1.822	2.894	3.842	4.809	3.659	3.273	1.154	0.459	0.110	0.105	0.003		0.033				
2003	4.819	3.324	2.401	3.516	3.757	2.245	1.743	0.749	0.423	0.207	0.024	0.004		0.026		0.016		
2004	4.722	3.217	3.000	3.430	3.605	2.358	1.490	0.572	0.311	0.113	0.106	0.005			0.003			
2005	0.037	1.264	1.723	3.226	2.716	2.107	1.321	0.473	0.263	0.155	0.028	0.064						
2006	6.705	5.126	2.126	3.172	2.692	1.936	1.847	1.129	0.177	0.130	0.012	0.023	0.004					
2007	26.051	2.543	3.567	3.118	4.005	2.557	1.703	1.258	0.456	0.123	0.026	0.014		0.005				
2008	13.880	2.399	1.815	1.733	1.573	1.015	0.763	0.425	0.230	0.099	0.026	0.023	0.025	0.000	0.000			
2009	2.032	3.973	1.945	2.898	3.289	1.738	0.812	0.471	0.558	0.199	0.033	0.065	0.002		0.001	0.002		
2010	1.300	5.701	2.689	3.141	2.522	1.978	0.681	0.364	0.465	0.248	0.120	0.052	0.023	0.006	0.002	0.004		
2011	0.518	3.795	3.527	2.746	3.011	2.018	1.544	0.421	0.355	0.149	0.094	0.019	0.060	0.013				
2012	0.098	3.650	2.315	3.724	2.026	1.343	0.913	0.541	0.256	0.109	0.124	0.049	0.019	0.024	0.006		0.003	
2013	0.583	5.142	3.306	1.857	1.960	1.510	0.952	0.695	0.451	0.216	0.088	0.089	0.062	0.004	0.006	0.002	0.006	
2014	17.884	6.474	4.500	3.324	2.337	3.135	1.714	1.202	0.698	0.509	0.098	0.087	0.082	0.007	0.007	0.025	0.013	0.015
2015	0.262	4.888	5.054	3.311	2.849	1.434	1.489	0.560	0.411	0.370	0.161	0.038	0.052	0.003	0.012		0.001	

2016	1.276	2.990	3.913	4.900	3.053	2.741	0.961	0.773	0.530	0.249	0.132	0.242	0.041	0.007	0.013		0.011	
2017	6.506	1.063	3.440	3.298	2.524	1.884	1.209	0.497	0.282	0.185	0.054	0.095	0.020	0.014				
2018	0.690	5.028	2.993	2.013	2.606	1.581	1.151	0.522	0.267	0.196	0.081	0.057	0.013	0.013		0.004	0.001	
2019	0.925	3.464	3.443	4.787	3.112	3.160	1.942	1.222	0.317	0.384	0.158	0.059	0.094	0.013	0.003	0.007	0.005	
2020	0.169	0.498	1.474	2.583	2.927	1.873	0.877	0.563	0.313	0.176	0.076	0.051		0.000	0.005	0.001		

Table A.1.2. CV on abundance indices

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17
1995	1.007	0.165	0.103	0.063	0.112	0.125	0.200	0.163	0.308	0.477	0.408		0.221					
1996	0.687	0.197	0.093	0.105	0.118	0.105	0.217	0.437	0.393	0.533	0.301							
1997	0.970	0.228	0.214	0.134	0.134	0.127	0.229	0.218	0.412	0.903	0.815	1.060						
1998	0.856	0.249	0.118	0.094	0.139	0.142	0.266	0.509	0.313	0.864	1.039		0.935		0.000	0.000		
1999		0.237	0.092	0.093	0.169	0.154	0.184	0.157	0.459	0.132	0.923		0.917					
2000	0.877	0.136	0.145	0.097	0.076	0.080	0.094	0.133	0.244	0.399	0.393	0.333		0.903				
2001	1.052	0.178	0.294	0.163	0.139	0.161	0.163	0.201	0.286	0.701	0.689	0.475	0.745	0.864				
2002	0.859	0.162	0.126	0.136	0.101	0.088	0.062	0.149	0.253	0.704	0.563	1.070		0.902				
2003	0.310	0.262	0.238	0.122	0.120	0.113	0.142	0.160	0.188	0.245	0.344	1.032		1.687			1.009	
2004	0.185	0.158	0.135	0.116	0.086	0.096	0.130	0.125	0.175	0.644	0.238	0.611			0.808			
2005	0.915	0.131	0.159	0.124	0.067	0.096	0.130	0.117	0.265	0.329	0.326	0.843						
2006	0.519	1.048	0.584	0.122	0.119	0.115	0.240	0.205	0.168	0.212	0.000	0.841	0.992					
2007	0.634	0.203	0.176	0.174	0.142	0.156	0.150	0.201	0.297	0.433	0.841	1.296		1.253				
2008	0.343	0.238	0.182	0.138	0.129	0.114	0.136	0.187	0.231	0.379	0.373	0.476	0.501	0.896	0.896			
2009	1.030	0.213	0.152	0.106	0.095	0.134	0.106	0.154	0.135	0.234	0.308	0.205	0.863		1.097	0.601		
2010	0.509	0.395	0.165	0.147	0.114	0.101	0.185	0.190	0.207	0.195	0.202	0.206	0.567	0.292	0.789	0.566		
2011	0.827	0.125	0.126	0.110	0.119	0.108	0.125	0.252	0.163	0.192	0.277	0.517	0.338	0.866				
2012	0.596	0.200	0.185	0.122	0.128	0.126	0.098	0.177	0.162	0.233	0.179	0.415	0.465	0.559	0.572		0.795	
2013	0.543	0.129	0.149	0.120	0.133	0.119	0.118	0.165	0.195	0.312	0.345	0.357	0.485	0.919	0.569	0.905	0.666	

2014	1.038	0.145	0.173	0.110	0.137	0.138	0.119	0.139	0.088	0.210	0.392	0.331	0.320	0.926	0.583	0.493	0.742	0.898
2015	0.706	0.151	0.097	0.089	0.068	0.086	0.096	0.142	0.140	0.130	0.271	0.360	0.427	0.804	0.554		1.010	
2016	0.564	0.281	0.117	0.103	0.074	0.115	0.141	0.160	0.195	0.347	0.304	0.452	0.431	0.399	0.526		2.264	
2017	0.635	0.326	0.151	0.119	0.151	0.133	0.166	0.206	0.257	0.421	0.246	0.410	0.522	0.746				
2018	0.413	0.175	0.157	0.155	0.148	0.143	0.071	0.173	0.554	0.141	0.345	0.428	0.603	0.503		0.515	1.528	
2019	0.232	0.137	0.172	0.117	0.093	0.083	0.101	0.100	0.194	0.232	0.235	0.207	0.631	0.518	1.053	1.170	1.069	
2020	0.827	0.320	0.134	0.158	0.111	0.133	0.141	0.220	0.299	0.466	0.527	0.433		1.148	1.151	1.011		

Table A.1.3. Biomass indices (kilotonnes)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17
1995	0.000	1.595	3.414	5.298	10.124	14.865	12.579	7.364	4.352	2.460	0.144		2.387					
1996	0.151	0.716	2.459	9.286	9.228	14.394	9.253	5.554	1.402	0.378	2.020							
1997	0.000	0.742	4.335	13.470	12.866	10.135	7.839	2.911	1.564	0.570	0.151	0.098						
1998	0.012	0.459	3.533	9.317	13.530	9.495	5.873	4.444	0.832	0.675	0.587		0.106		0.525	0.690		
1999		0.163	1.110	3.412	5.253	5.734	2.634	0.974	0.225	0.768	0.033		0.083					
2000	0.023	0.368	2.708	5.928	7.447	10.068	8.850	3.148	2.040	0.714	0.343	0.322		0.042				
2001	0.001	0.202	1.940	4.520	7.163	5.798	5.105	3.435	1.292	0.213	0.453	0.516	0.078	0.275				
2002	0.004	0.207	1.324	4.145	10.772	12.037	15.280	7.214	2.094	0.325	0.688	0.066		0.403				
2003	0.044	0.232	0.878	2.892	5.256	4.996	5.326	2.982	2.183	1.278	0.216	0.008		0.288		0.404		
2004	0.031	0.286	1.059	3.022	6.128	5.124	4.709	2.130	1.265	0.479	1.015	0.046		0.025				
2005	0.001	0.132	0.724	3.068	5.706	5.325	3.886	1.927	0.906	0.745	0.378	1.061						
2006	0.054	0.386	0.856	3.191	4.501	4.455	6.237	4.401	0.907	0.909	0.080	0.566	0.017					
2007	0.169	0.263	1.806	3.331	7.546	6.659	5.502	5.810	2.963	1.387	0.447	0.038		0.098				
2008	0.086	0.236	0.820	1.990	3.216	3.378	2.586	1.968	1.111	0.416	0.105	0.272	0.125	0.003	0.003			
2009	0.018	0.295	0.779	3.265	6.285	4.643	2.904	2.071	2.668	1.144	0.153	0.254	0.005		0.002	0.015		
2010	0.013	0.349	1.378	3.541	4.856	5.497	2.401	1.706	2.643	1.536	0.755	0.340	0.188	0.013	0.022	0.035		
2011	0.004	0.263	1.323	3.126	6.611	6.104	6.308	2.243	2.143	0.960	0.491	0.070	0.498	0.060				

2012	0.001	0.268	0.833	4.370	4.211	4.133	3.423	2.566	1.428	0.688	0.866	0.233	0.120	0.129	0.043		0.012	
2013	0.007	0.423	1.246	1.662	3.477	4.405	3.267	2.707	2.361	1.239	0.363	0.805	0.768	0.014	0.049	0.032	0.090	
2014	0.119	0.515	1.790	3.802	4.529	9.257	6.521	5.706	3.689	3.017	0.914	0.638	0.586	0.031	0.058	0.251	0.110	0.366
2015	0.001	0.372	2.001	3.673	5.571	4.061	5.054	2.655	2.078	2.252	0.970	0.335	0.257	0.031	0.118		0.015	
2016	0.009	0.219	1.432	6.363	6.806	9.072	3.621	3.389	3.349	1.915	0.851	2.480	0.209	0.032	0.111		0.095	
2017	0.024	0.084	1.782	3.530	5.057	5.635	4.825	2.249	1.998	1.168	0.337	0.902	0.102	0.213				
2018	0.004	0.257	1.541	2.341	5.458	5.214	4.334	2.817	1.598	1.406	0.572	0.233	0.065	0.113		0.079	0.007	
2019	0.008	0.230	1.264	5.626	6.381	9.351	7.683	5.747	1.634	2.419	1.091	0.477	1.047	0.127	0.027	0.082	0.038	
2020	0.001	0.043	0.800	3.417	5.736	6.224	3.032	3.247	1.916	1.357	0.469	0.365		0.001	0.036	0.013		

Table A.1.4. Length at age (cm)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17
1995	10.32	18.93	32.17	42.33	51.98	59.07	64.26	77.92	84.99	87.89	66.98		121.24					
1996	9.36	16.66	28.83	42.12	52.28	58.17	64.66	75.70	86.68	99.18	117.56							
1997	9.01	16.70	29.27	40.89	51.93	58.40	66.89	68.71	71.67	102.00	104.42	83.00						
1998	10.62	18.05	32.04	44.30	52.03	60.09	68.11	75.99	85.28	83.72	108.43		105.00		121.00	135.00		
1999		19.52	31.19	44.18	54.57	59.18	65.77	75.68	89.32	101.33	59.00		118.00					
2000	10.84	20.08	32.38	43.35	53.30	61.11	64.79	74.55	84.79	86.55	107.35	102.13		79.00				
2001	8.09	19.99	33.90	45.84	55.13	60.77	65.76	67.88	81.74	87.23	107.06	114.32	99.84	112.68				
2002	10.48	22.97	35.07	47.29	59.65	67.22	75.42	81.62	76.42	67.24	85.16	115.00		108.00				
2003	9.71	19.33	33.45	43.70	52.16	60.48	67.16	72.87	77.01	80.32	92.80	61.00		106.31		143.00		
2004	8.97	21.12	33.05	44.22	54.94	59.29	67.31	69.99	71.96	75.38	92.84	91.32		89.00				
2005	11.44	21.97	34.80	45.10	57.91	62.71	64.88	71.87	71.14	75.26	106.58	108.92						
2006	9.43	20.68	34.66	46.28	55.04	60.34	68.05	71.02	77.35	85.28	86.00	125.00	76.00					
2007	9.20	21.46	36.33	47.10	56.90	62.92	67.66	74.15	84.14	100.71	113.68	63.01		123.00				
2008	9.28	22.26	35.87	48.72	58.44	68.49	69.09	76.16	75.58	72.09	71.91	100.25	75.56	100.00	89.00			
2009	9.74	19.82	33.65	47.88	57.25	64.28	70.03	75.13	75.49	81.68	75.17	69.32	69.00		73.00	94.94		

2010	10.64	18.92	37.04	48.18	57.32	64.39	70.61	76.38	77.12	82.07	83.16	80.18	88.20	61.58	104.00	89.97			
2011	9.48	20.10	33.60	48.36	60.20	66.93	72.32	78.96	80.93	81.32	77.59	74.36	93.14	75.21					
2012	9.99	20.36	33.07	48.67	59.89	66.59	72.03	76.90	80.64	82.81	85.16	78.35	84.89	83.28	90.72		79.00		
2013	11.56	20.96	34.05	45.15	56.49	66.80	70.65	72.89	78.61	82.37	74.51	90.29	100.75	69.54	88.38	113.00	125.00		
2014	9.24	21.05	34.56	48.92	58.22	66.93	73.29	77.90	79.88	82.14	97.37	86.29	84.18	81.05	85.00	91.11	93.45	132.00	
2015	9.65	20.22	34.23	48.23	58.38	66.30	69.47	77.15	78.70	82.51	84.53	88.98	77.69	95.00	98.15		101.00		
2016	9.36	20.33	33.67	50.29	60.03	69.24	72.51	75.06	84.08	89.09	82.44	97.85	80.52	76.64	96.75		96.00		
2017	8.24	20.34	37.27	47.73	59.03	66.82	73.90	78.31	86.45	84.19	84.37	96.94	83.08	111.43					
2018	10.04	16.95	37.38	48.53	60.08	69.14	71.48	81.28	85.20	86.27	87.52	75.13	81.27	93.48		115.46	93.00		
2019	10.11	19.89	33.35	49.27	59.27	67.46	73.53	77.35	79.47	82.90	85.52	91.33	96.75	99.46	96.00	109.00	96.00		
2020	10.72	20.60	36.52	49.67	58.02	67.85	70.46	80.90	83.33	89.49	83.05	86.92		84.00	91.19	97.00			

Table A.1.5. Weight at age (gram)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17
1995	9	59	310	730	1405	2118	2754	4733	6494	7359	3297		17869					
1996	8	41	229	736	1385	1940	2702	4269	6627	10964	16213							
1997	8	42	240	665	1384	1932	2947	3175	4264	9600	11387	5140						
1998	10	53	332	869	1405	2229	3251	4432	5987	5884	16233		12260		19880	26100		
1999		64	278	828	1602	2059	2811	4567	7363	11535	2190		18340					
2000	12	70	320	798	1508	2329	2848	4424	6726	8851	12330	12873		4220				
2001	5	74	390	953	1648	2300	3116	3386	5983	7187	11653	17035	8460	15001				
2002	9	114	460	1077	2248	3295	4674	6289	4656	3785	7783	21980		12350				
2003	9	68	366	821	1397	2225	3056	3965	5110	6049	8878	2030		12562		25600		
2004	7	89	353	883	1702	2175	3158	3735	4004	4500	9936	9078		7130				
2005	16	105	419	954	2103	2529	2952	4081	3466	4745	14164	15482						
2006	8	84	424	1006	1673	2305	3380	3944	5151	6931	6530	25080	4220					
2007	7	103	508	1069	1889	2612	3238	4638	6497	11158	15891	2981		21100				

2008	6	100	453	1148	2046	3321	3392	4648	4869	4344	4077	10683	4730	9250	8520			
2009	10	75	401	1127	1911	2668	3569	4374	4764	5720	4819	3817	3084		3122	9672		
2010	10	64	507	1127	1927	2782	3545	4693	5714	6198	6336	6547	7962	2062	12154	7475		
2011	8	69	374	1136	2195	3020	4096	5365	6040	6395	5275	3933	8509	4558				
2012	8	74	359	1170	2084	3086	3748	4717	5530	6291	7000	5018	7205	6444	7052		4630	
2013	12	83	377	893	1773	2915	3432	3911	5284	5662	4216	8854	12081	4092	7780	16220	16300	
2014	7	80	399	1142	1935	2951	3804	4750	5282	5933	8898	7135	7062	4752	7920	9591	8282	24950
2015	8	76	395	1109	1956	2832	3397	4764	5073	6059	6231	8733	5427	10170	9930		11240	
2016	7	74	367	1295	2230	3307	3777	4399	6283	7696	6307	10446	5020	4413	9411		8325	
2017	4	83	519	1073	2006	2984	3990	4515	7030	6363	6225	9634	5089	14019				
2018	5	52	512	1169	2099	3305	3771	5424	5961	7088	7024	4283	5345	9367		15761	7350	
2019	9	67	367	1176	2049	2959	3958	4696	5085	6396	6804	8025	10306	9786	9585	12345	7040	
2020	8	91	538	1313	1958	3318	3457	5706	6190	7737	6667	7430		5165	7911	9765		

Table A.1.6. Abundance index, standard deviation (SD) and Coefficient of variation (CV) for sum of age 2+ fish

	Index_2plus	SD_2plus	CV_2plus
1995		40.034	2.670
1996		42.500	2.417
1997		56.823	5.676
1998		38.388	2.848
1999		15.451	1.184
2000		29.426	2.112
2001		19.437	2.667
2002		20.341	1.462
2003		15.111	1.527
2004		14.994	1.001
2005		12.075	0.766

2006		13.247		1.345		0.101
2007		16.833		1.429		0.085
2008		7.728		0.770		0.100
2009		12.012		0.886		0.074
2010		12.295		0.910		0.074
2011		13.958		0.921		0.066
2012		11.452		0.939		0.082
2013		11.204		0.992		0.089
2014		17.758		1.520		0.086
2015		15.744		0.858		0.054
2016		17.567		0.860		0.049
2017		13.501		1.118		0.083
2018		11.498		1.015		0.088
2019		18.706		1.279		0.068
2020		10.920		0.780		0.071

## 2.2 - Subarea A: North of 67°N

Table A.2.1. Abundance indices (millions)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17
1995	0.031	26.495	8.774	4.974	6.382	6.440	4.373	1.309	0.532	0.319	0.041		0.090					
1996	21.458	17.580	9.025	8.592	4.576	5.306	2.723	1.022	0.213	0.032	0.024							
1997	0.018	16.567	15.358	16.930	7.710	4.484	2.316	0.716	0.328	0.059	0.014	0.019						
1998	1.260	8.360	6.757	8.524	8.261	3.717	1.530	0.700	0.102	0.122	0.037		0.009		0.000	0.000		
1999		2.494	3.486	3.387	2.788	2.498	0.751	0.172	0.030	0.022	0.015		0.005					
2000	1.979	5.028	7.439	5.831	3.939	3.853	2.825	0.622	0.258	0.071	0.013	0.010		0.010				
2001	0.207	2.711	4.551	4.246	3.776	2.184	1.499	0.974	0.149	0.029	0.036	0.029	0.009	0.018				

2002	0.418	1.188	2.071	2.532	2.926	2.075	0.970	0.596	0.293	0.106	0.089	0.003		0.033				
2003	4.798	3.276	2.168	3.026	3.303	1.838	1.519	0.651	0.364	0.190	0.024	0.003		0.026			0.016	
2004	4.431	3.046	2.643	2.819	2.589	1.686	1.094	0.371	0.213	0.104	0.064	0.005			0.003			
2005	0.019	0.904	1.201	2.228	1.816	1.490	0.843	0.234	0.233	0.127	0.015	0.064						
2006	6.231	4.981	1.836	2.587	2.210	1.453	1.612	1.046	0.130	0.089	0.000	0.023	0.004					
2007	26.051	2.458	3.037	2.778	3.794	2.437	1.632	1.215	0.441	0.120	0.023	0.014		0.005				
2008	13.853	2.344	1.739	1.684	1.511	0.985	0.761	0.399	0.225	0.097	0.026	0.023	0.024	0.000	0.000			
2009	1.804	3.907	1.502	2.084	2.596	1.373	0.605	0.386	0.378	0.140	0.031	0.029	0.002		0.001	0.002		
2010	1.170	5.509	2.503	2.853	2.240	1.679	0.583	0.309	0.432	0.229	0.113	0.052	0.023	0.002	0.002	0.004		
2011	0.363	2.104	2.542	1.869	2.372	1.469	1.215	0.394	0.278	0.137	0.074	0.018	0.046	0.013				
2012	0.098	3.561	2.170	3.546	1.832	1.154	0.791	0.503	0.254	0.107	0.124	0.049	0.019	0.024	0.006		0.003	
2013	0.421	4.694	3.084	1.597	1.770	1.287	0.838	0.657	0.430	0.216	0.083	0.089	0.062	0.004	0.006	0.002	0.006	
2014	16.680	6.030	4.171	3.066	2.137	2.904	1.609	1.151	0.429	0.462	0.089	0.087	0.082	0.007	0.007	0.025	0.013	0.015
2015	0.262	3.421	3.122	2.465	1.802	1.017	1.128	0.477	0.363	0.303	0.158	0.038	0.052	0.003	0.012		0.001	
2016	1.272	2.921	3.341	3.667	2.349	2.308	0.841	0.669	0.452	0.222	0.115	0.123	0.041	0.004	0.013		0.011	
2017	6.506	1.018	3.289	3.202	2.335	1.764	1.122	0.450	0.256	0.181	0.054	0.095	0.020	0.014				
2018	0.680	4.977	2.847	1.837	2.376	1.246	0.946	0.494	0.246	0.136	0.081	0.057	0.013	0.013		0.004	0.001	
2019	0.305	2.607	2.992	3.724	2.221	2.149	1.272	0.656	0.212	0.262	0.106	0.040	0.092	0.013	0.003	0.007	0.005	
2020	0.162	0.475	1.039	1.743	2.204	1.329	0.674	0.363	0.246	0.074	0.075	0.047		0.000	0.004	0.001		

Table A.2.2. CV on abundance indices

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17
1995	1.007	0.168	0.130	0.090	0.124	0.136	0.211	0.194	0.395	0.505	0.412		0.328					
1996	0.700	0.198	0.111	0.154	0.168	0.141	0.259	0.538	0.394	0.516	1.565							
1997	0.970	0.239	0.246	0.160	0.156	0.142	0.250	0.259	0.470	0.903	0.815	1.060						
1998	0.860	0.259	0.186	0.117	0.161	0.161	0.311	0.690	0.401	0.867	1.039		0.935					
1999		0.243	0.105	0.114	0.198	0.173	0.229	0.193	0.472	0.402	0.923		0.917					

2000	0.877	0.143	0.164	0.124	0.095	0.090	0.103	0.151	0.291	0.489	0.894	0.901		0.903				
2001	1.052	0.179	0.313	0.182	0.159	0.186	0.176	0.209	0.406	0.719	0.689	0.475	0.745	0.864				
2002	0.859	0.248	0.174	0.205	0.160	0.136	0.147	0.228	0.363	0.723	0.665	1.070		0.902				
2003	0.311	0.266	0.263	0.140	0.137	0.137	0.161	0.182	0.218	0.262	0.346	0.694		1.687			1.009	
2004	0.196	0.166	0.153	0.141	0.118	0.127	0.171	0.172	0.253	0.694	0.330	0.611			0.808			
2005	0.774	0.181	0.228	0.177	0.099	0.136	0.204	0.228	0.298	0.400	0.606	0.843						
2006	0.559	1.079	0.677	0.149	0.145	0.153	0.275	0.221	0.228	0.311		0.841	0.992					
2007	0.634	0.209	0.201	0.194	0.149	0.163	0.156	0.207	0.307	0.446	0.969	1.296		1.253				
2008	0.343	0.244	0.189	0.142	0.134	0.118	0.136	0.198	0.236	0.386	0.373	0.481	0.521	0.896	0.896			
2009	1.160	0.217	0.197	0.146	0.118	0.168	0.139	0.183	0.192	0.314	0.246	0.442	0.863		1.097	0.601		
2010	0.563	0.408	0.177	0.162	0.128	0.119	0.216	0.223	0.223	0.210	0.215	0.207	0.567	1.171	0.789	0.566		
2011	1.075	0.218	0.170	0.159	0.146	0.147	0.155	0.268	0.205	0.195	0.347	0.510	0.443	0.866				
2012	0.598	0.205	0.197	0.129	0.141	0.147	0.113	0.190	0.163	0.238	0.179	0.415	0.465	0.559	0.572		0.795	
2013	0.681	0.140	0.160	0.140	0.147	0.139	0.134	0.174	0.204	0.312	0.365	0.357	0.485	0.919	0.569	0.905	0.666	
2014	1.112	0.155	0.186	0.119	0.150	0.149	0.127	0.146	0.143	0.231	0.433	0.331	0.320	0.926	0.583	0.493	0.742	0.898
2015	0.706	0.211	0.157	0.114	0.100	0.118	0.121	0.160	0.157	0.163	0.267	0.360	0.424	0.804	0.554		1.010	
2016	0.565	0.290	0.152	0.096	0.078	0.105	0.162	0.169	0.208	0.390	0.313	0.225	0.431	0.647	0.526		2.264	
2017	0.635	0.341	0.158	0.123	0.163	0.142	0.179	0.227	0.282	0.432	0.246	0.410	0.522	0.746				
2018	0.420	0.177	0.165	0.171	0.162	0.180	0.087	0.184	0.604	0.204	0.345	0.428	0.603	0.503		0.515	1.528	
2019	0.679	0.180	0.198	0.147	0.127	0.120	0.151	0.176	0.283	0.333	0.348	0.275	0.636	0.518	1.053	1.170	1.069	
2020	0.833	0.326	0.141	0.148	0.133	0.137	0.157	0.224	0.313	0.472	0.526	0.419		1.148	0.817	1.011		

Table A.2.3. Biomass indices (kilotonnes)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17
1995	0.000	1.541	2.472	3.574	8.904	13.457	11.950	6.114	3.126	2.269	0.143		1.576					
1996	0.146	0.709	1.947	5.779	6.165	10.249	7.685	4.485	1.401	0.368	0.474							
1997	0.000	0.677	3.790	11.041	10.756	8.589	6.793	2.140	1.152	0.570	0.151	0.098						

1998	0.012	0.410	1.751	7.160	11.622	8.458	4.866	3.312	0.527	0.674	0.587		0.106		0.000	0.000			
1999		0.155	0.947	2.695	4.204	4.894	2.062	0.733	0.217	0.144	0.033		0.083						
2000	0.023	0.347	2.398	4.816	6.154	9.109	7.947	2.654	1.529	0.406	0.111	0.088		0.042					
2001	0.001	0.199	1.778	3.973	6.276	5.075	4.503	3.240	0.679	0.205	0.453	0.516	0.078	0.275					
2002	0.004	0.105	0.737	2.351	4.653	4.935	3.358	2.646	1.109	0.289	0.553	0.066		0.403					
2003	0.044	0.228	0.783	2.488	4.716	4.167	4.746	2.687	2.016	1.232	0.216	0.006		0.288			0.404		
2004	0.029	0.269	0.896	2.463	4.253	3.623	3.504	1.407	1.012	0.418	0.320	0.046			0.025				
2005	0.000	0.089	0.526	1.947	3.135	3.291	2.133	0.860	0.813	0.706	0.129	1.061							
2006	0.050	0.368	0.695	2.558	3.643	3.235	5.626	4.119	0.575	0.714	0.000	0.566	0.017						
2007	0.169	0.239	1.469	2.959	7.057	6.266	5.159	5.471	2.811	1.347	0.400	0.038		0.098					
2008	0.086	0.225	0.742	1.869	2.978	3.285	2.581	1.807	1.097	0.402	0.105	0.272	0.123	0.003	0.003				
2009	0.014	0.288	0.537	2.152	4.877	3.708	2.308	1.786	1.960	0.763	0.150	0.164	0.005		0.002	0.015			
2010	0.013	0.329	1.270	3.110	4.191	4.605	2.075	1.445	2.195	1.434	0.723	0.340	0.188	0.004	0.022	0.035			
2011	0.003	0.125	1.024	2.183	5.403	4.579	4.488	2.024	1.561	0.855	0.388	0.067	0.360	0.060					
2012	0.001	0.259	0.774	4.060	3.700	3.344	2.920	2.374	1.422	0.657	0.866	0.233	0.120	0.129	0.043		0.012		
2013	0.005	0.399	1.184	1.469	3.215	3.913	2.883	2.593	2.101	1.239	0.350	0.805	0.768	0.014	0.049	0.032	0.090		
2014	0.107	0.480	1.497	3.446	4.057	8.515	5.938	5.346	2.390	2.572	0.845	0.638	0.586	0.031	0.058	0.251	0.110	0.366	
2015	0.001	0.252	1.275	2.748	3.860	3.036	4.249	2.297	1.916	1.795	0.962	0.335	0.254	0.031	0.118		0.015		
2016	0.009	0.213	1.151	4.034	4.465	7.685	3.293	3.123	2.677	1.632	0.793	1.262	0.209	0.019	0.111		0.095		
2017	0.024	0.080	1.655	3.379	4.591	5.205	4.488	2.104	1.811	1.131	0.337	0.902	0.102	0.213					
2018	0.004	0.255	1.494	2.026	4.962	3.986	3.552	2.648	1.448	1.163	0.572	0.233	0.065	0.113		0.079	0.007		
2019	0.002	0.163	1.112	4.212	4.414	6.413	4.854	3.382	1.272	1.623	0.840	0.392	1.042	0.127	0.027	0.082	0.038		
2020	0.001	0.041	0.384	1.749	4.414	3.853	2.271	1.786	1.643	0.652	0.466	0.342		0.001	0.029	0.013			

Table A.2.4. Length by age (cm)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17
1995	10.32	18.86	31.39	42.05	51.83	58.77	64.27	77.55	82.36	87.08	67.02		123.50					

1996	9.35	16.68	28.32	41.25	51.87	58.06	65.16	74.80	86.75	99.57	115.00												
1997	9.01	16.61	29.56	40.75	51.99	58.11	66.89	66.80	68.62	102.00	104.42	83.00											
1998	10.62	17.76	30.26	44.04	51.99	60.31	67.81	74.93	82.22	83.81	108.43		105.00										
1999		19.41	31.17	44.06	54.06	58.69	65.44	74.00	88.95	88.21	59.00		118.00										
2000	10.84	20.01	32.55	43.97	53.96	61.39	64.53	73.81	81.94	80.31	95.00	95.83		79.00									
2001	8.09	19.97	33.70	45.70	55.37	61.09	65.17	67.64	76.07	87.23	107.08	114.32	99.84	112.68									
2002	10.48	21.64	32.65	45.02	54.46	62.01	68.83	72.35	70.52	66.73	85.06	115.00		108.00									
2003	9.71	19.30	33.32	43.76	52.60	60.94	67.73	73.67	78.79	81.86	92.81	61.00		106.31							143.00		
2004	8.96	21.11	32.71	44.03	54.46	59.25	67.72	70.52	75.47	74.17	78.15	91.32			89.00								
2005	10.83	21.55	35.73	44.69	55.45	60.55	62.61	71.42	71.73	80.28	93.20	108.92											
2006	9.36	20.55	34.08	46.19	54.99	59.98	68.76	71.39	74.57	89.02		125.00	76.00										
2007	9.20	21.18	35.90	47.15	56.79	62.73	67.32	73.73	83.42	100.54	116.00	63.01		123.00									
2008	9.28	22.14	35.38	48.30	57.91	68.55	69.09	75.83	75.85	71.71	71.91	100.50	76.19	100.00	89.00								
2009	9.31	19.77	32.91	46.75	57.08	64.72	71.36	76.56	76.92	81.20	75.56	77.44	69.00		73.00	94.94							
2010	10.95	18.85	36.94	47.84	56.92	64.14	71.22	76.43	75.46	82.06	83.12	80.25	88.20	60.00	104.00	89.97							
2011	9.48	19.13	34.62	48.69	61.02	67.55	71.22	78.14	80.80	80.53	78.30	74.68	91.37	75.21									
2012	9.99	20.30	32.93	48.26	59.32	65.46	71.40	76.44	80.72	82.21	85.18	78.35	84.89	83.28	90.72		79.00						
2013	11.52	21.22	34.28	45.56	56.92	67.73	70.94	73.28	77.28	82.37	75.33	90.29	100.75	69.54	88.38	113.00	125.00						
2014	9.12	21.05	33.75	48.79	57.97	66.89	72.83	77.47	81.67	80.79	97.65	86.29	84.20	81.05	85.00	91.11	93.45	132.00					
2015	9.65	19.89	34.64	48.32	60.32	67.78	72.64	77.88	79.92	82.17	84.77	88.98	77.76	95.00	98.15		101.00						
2016	9.36	20.29	33.09	48.16	58.03	69.50	73.48	76.86	82.46	87.50	83.78	92.38	80.52	81.82	96.75		96.00						
2017	8.24	20.30	36.98	47.56	58.69	66.70	73.97	79.48	85.96	83.97	84.37	96.94	83.08	111.43									
2018	10.03	16.97	37.62	48.03	60.12	68.65	71.49	81.07	84.73	92.15	87.52	75.13	81.27	93.48		115.46	93.00						
2019	9.51	19.64	33.67	49.05	59.01	68.22	73.51	80.38	84.43	84.10	91.63	97.77	97.12	99.46	96.00	109.00	96.00						
2020	10.70	20.47	33.36	46.64	58.76	65.99	70.57	77.97	85.63	90.86	83.04	87.25		84.00	91.43	97.00							

Table A.2.5. Weight by age (gram)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16	Age 17
1995	9	58	282	719	1395	2091	2767	4693	5905	7211	3300		17490					
1996	8	41	216	672	1349	1939	2779	4223	6638	11146	20000							
1997	8	41	244	655	1393	1914	2921	2988	3768	9600	11387	5140						
1998	10	49	259	840	1406	2261	3173	4320	5275	5896	16233		12260					
1999		63	272	793	1508	1964	2759	4257	7262	6561	2190		18340					
2000	12	69	322	826	1561	2363	2811	4260	5977	6061	8736	9458		4220				
2001	5	74	377	933	1660	2320	2998	3338	4478	7193	11658	17035	8460	15001				
2002	9	88	357	918	1595	2377	3468	4415	3868	3588	8921	21980		12350				
2003	9	68	361	820	1427	2269	3127	4114	5493	6350	8881	2030		12562			25600	
2004	6	88	338	877	1646	2153	3197	3810	4656	4184	5102	9078		7130				
2005	14	99	436	878	1727	2205	2542	3666	3520	5562	8810	15482						
2006	8	83	400	989	1649	2231	3502	3992	4445	8004		25080	4220					
2007	7	97	486	1066	1865	2579	3168	4520	6363	11111	17480	2981		21100				
2008	6	97	427	1109	1971	3327	3393	4543	4921	4270	4077	10741	4962	9250	8520			
2009	8	74	357	1032	1878	2695	3803	4599	5146	5349	4886	5474	3084		3122	9672		
2010	11	63	502	1088	1872	2745	3586	4684	5096	6263	6448	6562	7962	2635	12154	7475		
2011	8	59	401	1165	2279	3109	3702	5163	5593	6174	5325	3982	8151	4558				
2012	8	73	355	1141	2026	2907	3690	4688	5549	6118	7004	5018	7205	6444	7052		4630	
2013	12	85	384	918	1817	3041	3438	3963	4926	5662	4340	8854	12081	4092	7780	16220	16300	
2014	7	80	359	1122	1894	2929	3690	4646	5562	5550	9029	7135	7066	4752	7920	9591	8282	24950
2015	8	73	406	1115	2145	2987	3774	4839	5299	5869	6281	8733	5443	10170	9930		11240	
2016	7	73	347	1101	1904	3327	3928	4689	5885	7273	6709	10371	5020	5773	9411		8325	
2017	4	83	504	1058	1969	2943	3997	4676	6985	6306	6225	9634	5089	14019				
2018	5	52	522	1109	2094	3206	3763	5391	5818	8438	7024	4283	5345	9367		15761	7350	
2019	8	62	372	1131	1984	2983	3815	5141	5908	6420	7801	9778	10405	9786	9585	12345	7040	
2020	8	91	368	1002	2001	2904	3374	4938	6718	8514	6665	7555		5165	7984	9765		

Table A.2.6. Abundance index, standard deviation (SD) and Coefficient of variation (CV) for sum of age 2+ fish

	Index_2plus	SD_2plus	CV_2plus
1995	33.395	2.667	0.080
1996	31.513	2.386	0.076
1997	47.938	5.599	0.117
1998	29.757	2.844	0.096
1999	13.154	1.183	0.090
2000	24.871	2.111	0.085
2001	17.500	2.666	0.152
2002	11.695	1.446	0.124
2003	13.128	1.526	0.116
2004	11.593	0.990	0.085
2005	8.253	0.766	0.093
2006	10.989	1.345	0.122
2007	15.494	1.420	0.092
2008	7.476	0.770	0.103
2009	9.128	0.883	0.097
2010	11.022	0.909	0.083
2011	10.425	0.917	0.088
2012	10.581	0.939	0.089
2013	10.131	0.993	0.098
2014	16.259	1.520	0.093
2015	10.942	0.857	0.078
2016	14.157	0.795	0.056
2017	12.782	1.118	0.087
2018	10.298	1.015	0.099
2019	13.753	1.271	0.092

2020		7.800		0.674		0.086
------	--	-------	--	-------	--	-------

## 2.3 - Subarea B: Between 65° and 67°N

Table A.3.1. Abundance indices (millions)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
1995	0.000	0.319	1.494	1.205	0.390	0.219	0.204	0.155	0.117	0.020	0.000		0.044				
1996	0.586	0.065	1.390	3.276	1.178	1.406	0.425	0.134	0.000	0.000	0.099						
1997	0.000	0.934	1.928	2.312	1.023	0.383	0.094	0.036	0.005	0.000	0.000	0.000					
1998	0.000	0.264	1.216	0.903	0.846	0.406	0.070	0.053	0.039	0.000	0.000		0.000		0.026	0.026	
1999		0.036	0.360	0.466	0.260	0.202	0.127	0.023	0.001	0.045	0.000		0.000				
2000	0.000	0.236	0.976	1.511	0.814	0.350	0.244	0.089	0.049	0.016	0.016	0.016		0.000			
2001	0.000	0.110	0.315	0.358	0.382	0.399	0.162	0.064	0.065	0.005	0.000	0.000	0.004	0.000			
2002	0.013	0.203	0.422	0.342	0.375	0.119	0.116	0.045	0.032	0.000	0.016	0.000		0.000			
2003	0.021	0.085	0.293	0.764	0.518	0.436	0.237	0.116	0.085	0.023	0.000	0.004		0.000		0.000	
2004	0.061	0.163	0.499	0.708	0.955	0.648	0.364	0.169	0.111	0.011	0.038	0.000		0.000			
2005	0.018	0.126	0.464	0.730	0.445	0.432	0.240	0.188	0.035	0.027	0.015	0.000					
2006	0.060	0.153	0.246	0.512	0.301	0.292	0.195	0.075	0.012	0.029	0.000	0.000	0.000				
2007	0.049	0.086	0.540	0.325	0.162	0.123	0.166	0.052	0.011	0.000	0.000	0.001		0.000			
2008	0.323	0.140	0.098	0.028	0.028	0.020	0.006	0.014	0.013	0.002	0.000	0.004	0.001	0.000	0.000		
2009	0.228	0.118	0.538	0.484	0.617	0.280	0.195	0.065	0.162	0.062	0.001	0.013	0.000		0.000	0.000	
2010	0.014	0.488	0.343	0.282	0.211	0.258	0.113	0.064	0.010	0.021	0.022	0.000	0.003	0.005	0.000	0.000	
2011	0.155	1.784	1.046	0.828	0.553	0.425	0.178	0.028	0.069	0.004	0.014	0.001	0.012	0.000			
2012	0.001	0.218	0.573	0.323	0.084	0.086	0.044	0.052	0.018	0.009	0.013	0.019	0.004	0.008	0.000	0.003	
2013	0.165	0.620	0.618	0.509	0.340	0.303	0.164	0.075	0.044	0.020	0.015	0.009	0.015	0.000	0.000	0.000	
2014	1.136	0.357	0.392	0.158	0.062	0.080	0.052	0.065	0.235	0.114	0.010	0.000	0.012	0.005	0.000	0.000	
2015	0.000	1.371	0.849	0.714	0.792	0.335	0.230	0.067	0.035	0.059	0.061	0.004	0.015	0.000	0.005	0.000	

2016	0.033	0.371	0.433	0.223	0.244	0.304	0.193	0.150	0.053	0.035	0.015	0.014	0.000	0.004	0.000		0.000
2017	0.000	0.055	0.209	0.120	0.053	0.123	0.054	0.059	0.022	0.013	0.013	0.009	0.005	0.002			
2018	0.010	0.289	0.158	0.173	0.094	0.093	0.109	0.068	0.042	0.021	0.012	0.004	0.002	0.006		0.000	0.000
2019	0.631	0.874	0.616	1.017	0.850	0.681	0.191	0.237	0.066	0.087	0.054	0.027	0.004	0.000	0.000	0.000	0.000
2020	0.007	0.026	0.300	0.469	0.677	0.254	0.191	0.092	0.078	0.028	0.003	0.009		0.000	0.005	0.000	

Table A.3.2. CV on abundance indices

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
1995		0.027	0.008	0.023	0.074	0.081	0.055	0.036	0.167	1.012	6.127		0.000				
1996	0.355	0.635	0.011	0.010	0.054	0.055	0.115	0.174			0.000						
1997		0.234	0.174	0.060	0.092	0.137	0.353	0.501	2.629								
1998		0.145	0.075	0.100	0.080	0.104	0.464	0.240	0.443	8.401					0.000	0.000	
1999		0.010	0.002	0.001	0.001	0.001	0.000	0.000	0.019	0.000							
2000		0.038	0.013	0.012	0.027	0.075	0.101	0.080	0.038	0.103	0.000	0.000					
2001		0.682	0.384	0.257	0.110	0.156	0.164	0.255	0.176	0.968			1.159				
2002	2.110	0.263	0.394	0.096	0.048	0.298	0.564	0.981	0.662		0.000						
2003	0.522	0.324	0.099	0.061	0.092	0.074	0.090	0.129	0.156	0.284		1.032					
2004	0.347	0.052	0.075	0.079	0.061	0.102	0.113	0.143	0.109	1.036	0.303						
2005	1.591	0.255	0.046	0.033	0.028	0.034	0.039	0.084	0.283	0.000	0.148						
2006	0.000	0.041	0.031	0.021	0.049	0.036	0.036	0.089	0.202	0.000							
2007	1.071	0.366	0.157	0.144	0.154	0.151	0.206	0.208	0.960			5.384					
2008	0.723	0.555	0.336	0.708	0.471	0.101	0.856	0.584	0.695	0.201		0.685	0.040				
2009	0.000	0.369	0.177	0.078	0.111	0.084	0.097	0.295	0.082	0.199	4.224	0.273					
2010	0.221	0.367	0.172	0.146	0.080	0.076	0.076	0.062	0.444	0.708	0.248	5.180	0.805	0.000			
2011	0.712	0.078	0.098	0.087	0.103	0.080	0.144	0.736	0.111	1.824	1.206	3.159	0.000				
2012	1.698	0.213	0.507	0.524	0.318	0.305	0.457	0.434	0.408	0.442	0.369	0.656	0.769	0.807		0.795	
2013	0.586	0.160	0.191	0.092	0.067	0.057	0.160	0.271	0.254	0.585	0.633	1.014	0.631				

2014	0.118	0.145	0.158	0.176	0.108	0.231	0.273	0.412	0.039	0.335	0.711		0.745	1.254			
2015		0.053	0.059	0.087	0.083	0.090	0.152	0.317	0.466	0.352	0.678	0.970	0.527		1.096		
2016	1.416	0.828	0.216	0.335	0.242	0.232	0.291	0.377	0.464	0.407	0.753	0.590		0.639			
2017		0.637	0.330	0.262	0.206	0.279	0.349	0.384	0.529	0.419	0.619	0.531	1.049	0.936			
2018	0.638	0.533	0.167	0.147	0.170	0.367	0.506	0.648	0.594	0.625	0.626	0.852	0.684	1.026			
2019	0.056	0.097	0.145	0.075	0.077	0.076	0.192	0.108	0.183	0.117	0.097	0.195	1.579				
2020	3.801	0.853	0.400	0.222	0.162	0.337	0.310	0.350	0.471	0.519	1.056	1.242			1.369		

Table A.3.3. Biomass indices (kilotonnes)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
1995	0.000	0.029	0.528	0.814	0.573	0.469	0.517	0.822	1.037	0.188	0.001		0.811				
1996	0.005	0.001	0.404	2.548	1.454	2.479	0.939	0.622	0.000	0.000	1.546						
1997	0.000	0.053	0.377	1.398	1.149	0.663	0.257	0.113	0.008	0.000	0.000	0.000					
1998	0.000	0.034	0.375	0.753	1.019	0.735	0.197	0.242	0.305	0.001	0.000		0.000		0.525	0.690	
1999		0.004	0.088	0.302	0.306	0.384	0.325	0.107	0.008	0.624	0.000		0.000				
2000	0.000	0.022	0.290	1.016	0.966	0.688	0.779	0.495	0.511	0.308	0.232	0.233		0.000			
2001	0.000	0.010	0.097	0.234	0.421	0.519	0.577	0.190	0.499	0.037	0.000	0.000	0.050	0.000			
2002	0.000	0.019	0.117	0.437	0.880	0.217	0.194	0.076	0.088	0.000	0.135	0.000		0.000			
2003	0.000	0.006	0.104	0.466	0.540	0.775	0.590	0.321	0.283	0.054	0.000	0.008		0.000		0.000	
2004	0.000	0.016	0.170	0.517	1.402	1.127	0.967	0.495	0.284	0.034	0.682	0.000		0.000			
2005	0.000	0.009	0.157	0.475	0.687	0.987	0.663	0.566	0.101	0.040	0.278	0.000					
2006	0.001	0.019	0.124	0.500	0.400	0.516	0.453	0.174	0.052	0.133	0.000	0.000	0.000				
2007	0.000	0.011	0.254	0.258	0.229	0.255	0.409	0.157	0.071	0.000	0.000	0.001		0.000			
2008	0.003	0.018	0.044	0.040	0.066	0.053	0.021	0.056	0.072	0.015	0.000	0.026	0.003	0.000	0.000		
2009	0.004	0.009	0.192	0.577	1.041	0.696	0.529	0.167	0.593	0.386	0.003	0.026	0.000		0.000	0.000	
2010	0.000	0.030	0.121	0.229	0.389	0.606	0.308	0.289	0.024	0.059	0.109	0.000	0.006	0.009	0.000	0.000	
2011	0.001	0.146	0.298	0.783	0.895	1.020	0.541	0.093	0.560	0.010	0.046	0.003	0.119	0.000			

2012	0.000	0.021	0.192	0.181	0.104	0.183	0.105	0.165	0.078	0.057	0.056	0.096	0.016	0.036	0.000		0.012
2013	0.002	0.037	0.147	0.321	0.393	0.675	0.400	0.231	0.199	0.085	0.038	0.031	0.061	0.000	0.000	0.000	0.000
2014	0.012	0.030	0.232	0.114	0.061	0.157	0.173	0.251	1.117	0.437	0.041	0.000	0.043	0.015	0.000	0.000	0.000
2015	0.000	0.106	0.233	0.759	1.233	0.793	0.522	0.207	0.127	0.460	0.228	0.019	0.066	0.000	0.036		0.000
2016	0.000	0.031	0.133	0.197	0.331	0.701	0.550	0.462	0.251	0.357	0.072	0.086	0.000	0.012	0.000		0.000
2017	0.000	0.005	0.073	0.104	0.101	0.309	0.173	0.190	0.084	0.077	0.058	0.040	0.020	0.024			
2018	0.000	0.015	0.047	0.271	0.155	0.249	0.347	0.265	0.225	0.097	0.060	0.015	0.019	0.037		0.000	0.000
2019	0.006	0.069	0.184	1.140	1.734	1.720	0.582	0.841	0.199	0.370	0.251	0.143	0.015	0.000	0.000	0.000	0.000
2020	0.000	0.002	0.118	0.505	1.165	0.630	0.605	0.420	0.328	0.126	0.013	0.056		0.000	0.025	0.000	

Table A.3.4. Mean length (cm)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
1995		21.94	33.66	41.58	52.72	60.33	64.29	80.92	94.71	99.01	60.00		117.00				
1996	9.77	13.00	31.63	42.75	51.16	57.11	61.34	80.03			118.00						
1997		18.27	27.13	39.84	49.63	57.20	64.53	75.17	54.90								
1998		24.38	31.97	43.56	49.98	57.23	65.97	77.02	93.29	57.00					121.00	135.00	
1999		23.93	29.19	40.33	49.29	57.53	64.02	79.87	97.41	107.50							
2000		21.57	31.06	40.66	49.45	58.17	67.45	79.64	98.84	109.64	114.00	105.00					
2001		21.77	31.73	41.09	48.33	51.45	67.05	65.76	90.48	89.36			117.00				
2002	10.00	21.90	30.75	48.87	60.98	55.74	53.59	54.19	64.97		97.00						
2003	10.00	19.94	32.96	39.46	47.12	56.05	62.67	65.60	69.22	62.20		61.00					
2004	8.77	21.73	32.25	42.14	51.94	55.44	63.72	65.75	64.19	72.07	119.47						
2005	14.00	20.14	31.65	40.05	52.32	60.57	63.73	65.16	65.74	55.00	118.27						
2006	11.67	23.74	36.59	45.65	51.04	55.71	61.58	62.43	77.27	75.96							
2007	9.94	23.46	35.55	43.44	51.65	58.62	62.02	67.97	89.28			57.00					
2008	10.73	25.27	36.20	53.57	62.11	63.98	69.17	71.80	79.42	88.53		88.78	69.99				
2009	12.11	20.79	32.27	48.34	54.21	61.43	64.91	63.54	71.00	81.67	59.00	59.22					

2010	12.04	19.15	32.56	42.28	55.82	60.52	64.49	74.88	61.81	66.77	81.14	58.00	60.00	62.00				
2011	9.38	21.22	30.50	45.76	54.64	62.29	66.89	72.58	83.47	61.36	69.07	64.00	102.00					
2012	10.16	22.09	32.88	39.24	51.02	59.62	63.48	67.44	74.07	83.81	74.30	78.35	73.00	75.00				79.00
2013	11.77	19.05	30.12	40.89	49.69	62.10	64.29	68.26	76.04	76.13	64.04	69.00	74.35					
2014	10.74	21.22	38.24	42.25	47.29	58.78	70.43	73.78	76.04	74.19	81.04		72.99	75.00				
2015		20.63	30.34	47.34	54.38	61.30	60.63	68.91	71.92	88.58	74.42	77.00	76.03		95.00			
2016	9.90	21.41	32.22	42.52	51.41	60.40	65.69	66.74	77.77	96.51	75.68	83.88		73.00				
2017		22.31	33.68	44.75	57.05	63.09	69.05	68.58	73.69	85.70	78.01	75.94	81.30	99.00				
2018	10.43	17.57	32.07	50.93	54.46	63.45	66.72	72.57	77.50	77.15	78.02	74.66	98.00	80.03				
2019	10.55	20.66	30.44	48.03	58.75	62.90	66.59	71.27	67.22	73.38	73.15	82.61	76.59					
2020	11.37	22.57	34.54	46.72	54.72	62.14	66.55	75.63	75.15	77.74	79.23	82.83			83.00			

Table A.3.5. Mean weight (gram)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
1995		91	354	676	1470	2148	2538	5295	8844	9406	2400		18580				
1996	8	17	291	778	1233	1763	2214	4688			15560						
1997		57	198	604	1124	1733	2787	3282	1506								
1998		128	309	835	1204	1809	2823	4634	7894	1655					19880	26100	
1999		115	245	648	1175	1906	2556	4722	9728	13870							
2000		92	297	672	1186	1962	3185	5549	10436	18769	14266	14380					
2001		96	316	670	1106	1305	3618	2991	7694	7230			12000				
2002	8	93	281	1286	2350	1887	1755	1800	2999		8358						
2003	9	67	354	610	1045	1780	2491	2787	3325	2370		2030					
2004	6	100	341	732	1468	1741	2663	2936	2558	3377	18589						
2005	25	75	340	652	1545	2286	2761	3002	2942	1456	18343						
2006	13	125	506	976	1329	1769	2318	2308	4444	4608							
2007	7	134	474	796	1421	2083	2516	3047	6673		1648						

2008	10	165	456	1543	2445	2691	3419	4277	5767	7574		6239	2733					
2009	16	76	361	1194	1697	2487	2720	2575	3664	6374	2096	2034						
2010	14	62	355	822	1852	2355	2731	4539	2398	2842	4951	2128	2285	1910				
2011	8	82	286	947	1620	2404	3064	4232	8231	2328	3590	2440	10116					
2012	9	96	338	581	1287	2206	2548	3215	4206	7036	4248	4862	3745	4325				4630
2013	13	60	241	631	1160	2225	2436	3071	4535	4334	2569	3425	4214					
2014	11	85	601	725	982	2003	3272	4015	4751	3973	4625		3634	3340				
2015		77	275	1061	1556	2373	2271	3186	3614	8353	3497	4310	4380		7785			
2016	9	87	307	860	1360	2292	2796	3035	4734	10174	4596	5961		3425				
2017		89	363	880	1901	2484	3206	3257	3827	6124	4306	4305	4379	10610				
2018	5	48	296	1587	1643	2592	3083	3993	5141	4668	4787	3684	8695	5988				
2019	10	79	299	1121	2042	2529	3058	3557	3014	4284	4647	5383	4261					
2020	9	86	394	1073	1720	2512	3123	4707	4240	4690	3524	6025			5434			

Table A.3.6. Abundance index, standard deviation (SD) and Coefficient of variation (CV) for sum of age 2+ fish

	Index_2plus	SD_2plus	CV_2plus
1995		3.848	0.030
1996		7.908	0.072
1997		5.782	0.365
1998		3.587	0.104
1999		1.483	0.001
2000		4.082	0.033
2001		1.753	0.309
2002		1.467	0.289
2003		2.474	0.049
2004		3.503	0.067
2005		2.576	0.025

2006	1.662	0.022	0.013
2007	1.379	0.090	0.066
2008	0.214	0.060	0.279
2009	2.416	0.197	0.081
2010	1.331	0.108	0.081
2011	3.157	0.150	0.048
2012	1.238	0.565	0.457
2013	2.113	0.138	0.065
2014	1.184	0.068	0.057
2015	3.167	0.147	0.046
2016	1.668	0.344	0.206
2017	0.682	0.189	0.277
2018	0.783	0.233	0.298
2019	3.830	0.106	0.028
2020	2.107	0.289	0.137

## 2.4 - Subarea C: Between 62°N and 65°N

Table A.4.1. Abundance index (millions)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
1995	0.000	0.147	0.748	1.075	0.435	0.364	0.042	0.097	0.029	0.001	0.000		0.000				
1996	0.004	0.118	0.333	0.816	0.966	0.761	0.249	0.113	0.000	0.002	0.000						
1997	0.000	0.223	0.621	1.083	0.556	0.376	0.242	0.167	0.059	0.000	0.000	0.000					
1998	0.009	0.089	2.701	1.303	0.520	0.115	0.206	0.198	0.000	0.000	0.000		0.000		0.000	0.000	
1999		0.031	0.145	0.259	0.235	0.094	0.062	0.019	0.000	0.000	0.000		0.000				
2000	0.000	0.000	0.053	0.085	0.182	0.117	0.036	0.000	0.000	0.000	0.000	0.000		0.000			
2001	0.000	0.102	0.350	0.462	0.379	0.288	0.137	0.052	0.030	0.005	0.000	0.000	0.004	0.000			

2002	0.013	0.604	0.930	0.969	1.552	1.570	2.387	0.604	0.167	0.004	0.000	0.000		0.000			
2003	0.000	0.037	0.063	0.314	0.146	0.096	0.044	0.027	0.031	0.010	0.000	0.003		0.000			0.000
2004	0.230	0.039	0.206	0.264	0.390	0.233	0.074	0.049	0.013	0.014	0.003	0.000		0.000			
2005	0.000	0.235	0.091	0.319	0.505	0.256	0.254	0.060	0.005	0.000	0.002	0.000					
2006	0.415	0.008	0.064	0.072	0.211	0.238	0.065	0.024	0.035	0.012	0.012	0.000	0.000				
2007	0.049	0.082	0.152	0.097	0.103	0.099	0.163	0.066	0.024	0.004	0.004	0.001		0.000			
2008	0.350	0.129	0.124	0.053	0.054	0.011	0.005	0.025	0.008	0.000	0.000	0.003	0.000	0.000	0.000		
2009	0.000	0.063	0.241	0.446	0.258	0.093	0.019	0.041	0.032	0.005	0.000	0.023	0.000		0.000	0.000	
2010	0.118	0.397	0.181	0.221	0.153	0.161	0.026	0.009	0.042	0.025	0.015	0.000	0.003	0.000	0.000	0.000	
2011	0.000	0.094	0.102	0.131	0.120	0.134	0.164	0.036	0.008	0.008	0.032	0.000	0.002	0.000			
2012	0.000	0.130	0.440	0.381	0.187	0.181	0.151	0.083	0.020	0.007	0.013	0.019	0.004	0.008	0.000		0.003
2013	0.003	0.172	0.396	0.249	0.150	0.095	0.108	0.052	0.065	0.020	0.010	0.009	0.015	0.000	0.000	0.000	
2014	0.116	0.425	0.205	0.180	0.173	0.250	0.152	0.101	0.106	0.161	0.019	0.000	0.012	0.005	0.000	0.000	
2015	0.000	0.149	1.109	0.183	0.338	0.094	0.200	0.034	0.053	0.058	0.058	0.004	0.015	0.000	0.005		0.000
2016	0.030	0.301	0.341	1.122	0.557	0.336	0.103	0.080	0.104	0.023	0.032	0.133	0.000	0.000	0.000		0.000
2017	0.000	0.098	0.209	0.086	0.176	0.147	0.112	0.085	0.042	0.009	0.013	0.009	0.005	0.002			
2018	0.000	0.238	0.086	0.094	0.181	0.330	0.265	0.073	0.056	0.081	0.012	0.004	0.002	0.006		0.000	0.000
2019	0.011	0.027	0.166	0.136	0.083	0.372	0.489	0.393	0.067	0.055	0.013	0.011	0.003	0.000	0.000	0.000	0.000
2020	0.000	0.003	0.277	0.554	0.137	0.403	0.113	0.136	0.011	0.101	0.003	0.005		0.000	0.003	0.000	

Table A.4.2. CV on abundance indices

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
1995		0.139	0.050	0.040	0.094	0.103	0.707	0.112	0.455	5.064							
1996	1.121	0.313	0.202	0.129	0.116	0.160	0.440	0.349	5.506	2.545							
1997		0.342	0.143	0.052	0.122	0.151	0.287	0.258	0.502								
1998	0.898	0.073	0.000	0.000	0.000	0.000	0.000	0.000									
1999		0.289	0.080	0.052	0.057	0.122	0.154	0.294									

2000			0.000	0.000	0.000	0.000	0.000											
2001		0.737	0.345	0.200	0.116	0.222	0.244	0.418	0.494	0.903	9.960		1.159					
2002	2.110	0.083	0.172	0.009	0.053	0.109	0.069	0.185	0.315	1.143								
2003		0.741	0.325	0.108	0.205	0.188	0.222	0.273	0.370	0.452	2.311	0.694						
2004	0.141	0.206	0.155	0.188	0.083	0.201	0.149	0.167	0.475	0.558	1.490							
2005		0.000	0.030	0.022	0.012	0.039	0.013	0.060	0.741		0.918							
2006	0.000	0.617	0.044	0.000	0.037	0.019	0.060	0.096	0.000	0.000	0.000							
2007	1.071	0.393	0.417	0.222	0.122	0.136	0.205	0.148	0.417	0.000	0.000	5.384						
2008	0.668	0.600	0.267	0.373	0.247	0.124	1.112	0.318	1.139			0.724						
2009		0.689	0.396	0.078	0.229	0.059	0.234	0.261	0.153	0.991	5.898	0.000						
2010	0.149	0.451	0.325	0.186	0.099	0.115	0.291	0.406	0.084	0.588	0.368		0.805					
2011	0.669	0.833	0.603	0.270	0.148	0.069	0.064	0.527	0.142	0.012	0.487	2.036	0.004	4.559				
2012		0.358	0.661	0.444	0.141	0.145	0.132	0.274	0.371	0.587	0.371	0.656	0.769	0.807				0.795
2013	2.824	0.527	0.296	0.192	0.150	0.182	0.240	0.388	0.172	0.585	0.956	1.014	0.631					
2014	0.666	0.158	0.360	0.152	0.043	0.075	0.096	0.269	0.086	0.238	0.365		0.740	1.254				
2015		0.119	0.025	0.166	0.070	0.068	0.091	0.202	0.206	0.220	0.691	0.970	0.504		1.096			
2016	1.529	1.024	0.631	0.341	0.191	0.388	0.505	0.557	0.566	0.581	0.616	0.757						
2017		0.359	0.330	0.367	0.061	0.235	0.170	0.264	0.271	0.629	0.619	0.531	1.049	0.936				
2018		0.644	0.344	0.277	0.083	0.105	0.208	0.617	0.440	0.159	0.626	0.852	0.684	1.026				
2019	0.907	0.907	0.426	0.161	0.185	0.052	0.035	0.039	0.130	0.134	0.538	0.550	0.930					
2020		1.741	0.533	0.392	0.496	0.294	0.562	0.607	0.602	0.736	1.003	1.036			0.977			

Table A.4.3. Biomass indices (kilotonnes)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
1995	0.000	0.025	0.414	0.910	0.647	0.940	0.112	0.428	0.189	0.002	0.000		0.000				
1996	0.000	0.006	0.110	0.982	1.667	1.735	0.652	0.447	0.001	0.010	0.000						
1997	0.000	0.012	0.168	1.030	0.961	0.883	0.790	0.658	0.403	0.000	0.000	0.000					

1998	0.000	0.015	1.406	1.404	0.889	0.302	0.809	0.890	0.000	0.000	0.000		0.000		0.000	0.000	
1999		0.004	0.075	0.415	0.743	0.455	0.247	0.134	0.000	0.000	0.000		0.000				
2000	0.000	0.000	0.020	0.096	0.328	0.270	0.125	0.000	0.000	0.000	0.000		0.000				
2001	0.000	0.009	0.167	0.455	0.595	0.593	0.305	0.158	0.250	0.036	0.000	0.000	0.050	0.000			
2002	0.000	0.094	0.602	1.357	5.275	7.009	12.038	4.644	0.964	0.036	0.000	0.000		0.000			
2003	0.000	0.002	0.011	0.136	0.141	0.170	0.110	0.070	0.143	0.036	0.001	0.006		0.000			0.000
2004	0.002	0.002	0.055	0.196	0.697	0.545	0.298	0.256	0.030	0.084	0.013	0.000			0.000		
2005	0.000	0.034	0.048	0.663	1.934	1.134	1.108	0.527	0.008	0.000	0.030	0.000					
2006	0.003	0.001	0.042	0.133	0.486	0.761	0.219	0.133	0.280	0.062	0.080	0.000	0.000				
2007	0.000	0.017	0.107	0.162	0.306	0.319	0.518	0.372	0.214	0.041	0.047	0.001		0.000			
2008	0.003	0.012	0.097	0.115	0.201	0.042	0.017	0.152	0.058	0.000	0.000	0.026	0.000	0.000	0.000	0.000	
2009	0.000	0.004	0.117	0.582	0.483	0.250	0.084	0.178	0.155	0.013	0.001	0.064	0.000		0.000	0.000	
2010	0.001	0.028	0.045	0.279	0.341	0.458	0.109	0.028	0.470	0.119	0.076	0.000	0.006	0.000	0.000	0.000	
2011	0.000	0.007	0.028	0.188	0.339	0.517	1.286	0.233	0.021	0.095	0.145	0.000	0.019	0.000			
2012	0.000	0.012	0.144	0.396	0.477	0.723	0.566	0.330	0.083	0.026	0.056	0.096	0.016	0.036	0.000		0.012
2013	0.000	0.014	0.085	0.128	0.132	0.229	0.415	0.182	0.459	0.085	0.025	0.031	0.061	0.000	0.000	0.000	
2014	0.001	0.031	0.102	0.278	0.445	0.762	0.742	0.517	0.430	0.882	0.109	0.000	0.043	0.015	0.000	0.000	
2015	0.000	0.016	0.496	0.180	0.537	0.236	0.506	0.203	0.221	0.237	0.220	0.019	0.064	0.000	0.036		0.000
2016	0.000	0.024	0.197	2.195	2.116	1.234	0.350	0.299	0.795	0.217	0.130	1.304	0.000	0.000	0.000	0.000	
2017	0.000	0.008	0.123	0.117	0.440	0.509	0.400	0.263	0.255	0.041	0.058	0.040	0.020	0.024			
2018	0.000	0.013	0.025	0.097	0.426	1.272	1.005	0.326	0.348	0.340	0.060	0.015	0.019	0.037		0.000	0.000
2019	0.000	0.002	0.032	0.309	0.281	1.281	2.282	1.752	0.294	0.495	0.057	0.077	0.012	0.000	0.000	0.000	0.000
2020	0.000	0.000	0.344	1.266	0.307	2.021	0.467	1.147	0.055	0.733	0.010	0.033		0.000	0.018	0.000	

Table A.4.4. Length by age (cm)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
1995		26.44	38.14	44.42	53.51	63.65	64.79	78.31	88.80	65.00							

1996	8.00	16.98	30.91	48.14	55.40	60.93	64.48	76.20	65.00	87.00										
1997		16.88	29.03	45.38	55.34	63.09	67.60	75.88	90.63											
1998	10.78	25.15	36.44	46.50	55.90	62.52	71.00	77.88												
1999		23.52	36.75	52.54	66.26	75.73	73.44	86.23												
2000			33.79	49.42	56.33	60.59	67.12													
2001		21.79	35.36	44.63	52.72	57.48	59.25	64.73	90.77	90.74	69.00		117.00							
2002	10.00	25.10	39.62	52.40	68.57	73.74	77.28	88.77	85.79	89.00										
2003		17.33	27.22	35.36	45.55	54.61	63.06	61.49	75.71	71.34	88.00	61.00								
2004	9.30	16.31	28.85	41.21	54.47	58.08	70.35	76.20	63.76	86.18	72.27									
2005		24.71	37.06	58.01	70.61	75.43	72.50	93.44	57.50		108.00									
2006	9.93	21.95	40.86	53.85	59.97	66.41	69.09	71.73	87.55	82.00	86.00									
2007	9.94	26.64	39.20	53.10	63.56	65.87	66.06	77.01	99.99	102.00	110.00	57.00								
2008	9.81	22.77	42.89	60.65	70.68	71.54	71.69	82.27	100.17			99.00								
2009		20.15	35.27	49.79	55.73	65.44	78.15	76.08	75.71	62.77	66.00	65.00								
2010	8.22	19.64	28.89	47.25	58.83	64.62	71.71	67.04	90.71	80.03	79.43		60.00							
2011	9.10	20.22	33.14	53.10	64.73	73.65	84.34	87.10	64.73	103.35	73.70	67.26	82.97	74.00						
2012		21.75	32.39	47.37	63.96	71.35	73.06	74.45	73.64	71.31	74.42	78.35	73.00	75.00					79.00	
2013	11.72	20.98	29.72	38.97	46.18	64.47	72.53	72.98	85.58	76.13	64.02	69.00	74.35							
2014	10.89	21.03	34.50	51.93	63.01	66.43	77.65	79.30	74.98	80.31	86.90		72.94	75.00						
2015		23.27	35.78	46.08	53.44	64.92	61.78	80.40	73.82	75.44	74.67	77.00	76.22		95.00					
2016	10.00	21.11	37.34	57.67	70.65	71.52	69.98	69.99	87.03	96.91	74.71	97.06								
2017		21.74	38.28	50.20	63.62	68.17	71.30	67.80	85.38	81.71	78.01	75.94	81.30	99.00						
2018		19.07	32.02	47.30	61.82	72.74	71.57	78.07	84.51	74.02	78.02	74.66	98.00	80.03						
2019	14.17	19.07	28.08	55.60	66.71	69.89	76.19	74.94	74.76	91.90	77.99	92.20	76.83							
2020		18.35	47.79	57.70	60.23	76.12	71.63	88.75	79.70	87.56	79.00	83.00		83.00						

Table A.4.5. Weight by age (gram)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
1995		171	555	846	1489	2590	2748	4417	6905	2996							
1996	5	59	330	1212	1730	2284	2632	3977	2970	5220							
1997		58	275	952	1730	2359	3239	3982	7487								
1998	10	167	520	1078	1710	2620	3931	4494									
1999		120	517	1605	3163	4878	3958	7302									
2000			376	1134	1799	2305	3429										
2001		96	499	1005	1580	2089	2210	3051	9206	7802	2808		12000				
2002	8	156	658	1401	3400	4481	5053	7831	5977	8640							
2003		41	166	436	977	1800	2509	2644	4551	4036	7366	2030					
2004	9	44	267	762	1796	2407	4078	5302	2476	6949	3748						
2005		144	523	2081	3833	4438	4361	8869	1676		12140						
2006	8	81	652	1839	2300	3198	3350	5670	7957	5035	6530						
2007	7	244	844	1736	3018	3266	3246	5671	9142	11610	13370	1648					
2008	8	105	822	2259	3862	3886	3739	6215	10013			7675					
2009		71	540	1313	1932	2698	4495	4400	4936	2729	2656	2730					
2010	4	73	252	1301	2245	2869	4313	3162	11121	5156	5031		2285				
2011	7	66	386	1511	2859	3864	7854	7474	2569	11848	4739	2910	7858	3888			
2012		93	336	1191	2578	4054	3779	4088	4066	3716	4266	4862	3745	4325			4630
2013	12	78	217	515	889	2398	3969	3546	7124	4334	2499	3425	4214				
2014	15	73	539	1565	2571	3057	4901	5286	4051	5676	6016		3631	3340			
2015		112	448	1002	1592	2534	2532	6045	4153	4084	3532	4310	4410		7785		
2016	9	82	494	1952	3856	3623	3407	3576	7134	9101	4101	8575					
2017		87	637	1473	2499	3535	3579	3084	6189	4555	4306	4305	4379	10610			
2018		55	301	1078	2353	3858	3817	4955	6558	4188	4787	3684	8695	5988			
2019	23	61	190	2317	3433	3443	4671	4456	4388	9163	4210	7453	4298				
2020		48	1169	2194	2153	4935	3913	7895	5101	6869	3386	6140			5434		

Table A.4.6. Abundance index, standard deviation (SD) and Coefficient of variation (CV) for sum of age 2+ fish

	Index_2plus	SD_2plus	CV_2plus
1995	2.791	0.052	0.019
1996	3.241	0.168	0.052
1997	3.103	0.141	0.045
1998	5.044	0.000	0.000
1999	0.815	0.020	0.025
2000	0.474	0.000	0.000
2001	1.707	0.312	0.183
2002	8.183	0.353	0.043
2003	0.735	0.038	0.052
2004	1.246	0.038	0.031
2005	1.491	0.013	0.009
2006	0.735	0.018	0.025
2007	0.712	0.067	0.094
2008	0.282	0.060	0.211
2009	1.158	0.195	0.169
2010	0.835	0.108	0.130
2011	0.739	0.139	0.188
2012	1.498	0.565	0.377
2013	1.169	0.138	0.118
2014	1.363	0.079	0.058
2015	2.150	0.135	0.063
2016	2.832	0.597	0.211
2017	0.895	0.189	0.211
2018	1.191	0.236	0.198
2019	1.788	0.082	0.046

2020		1.743		0.341		0.196
------	--	-------	--	-------	--	-------

## 11 - Appendix B Swept area abundance indices

### 3.1 - Total area

Table B.1.1. Abundance indices (millions)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	6.253	5.298	3.531	4.329	5.087	3.391	2.719	0.999	0.577	0.271	0.043	0.002		0.016		0.010	
2004	5.077	2.987	2.495	2.885	3.852	2.109	1.079	0.560	0.238	0.054	0.072	0.006			0.006		
2005	0.030	0.906	1.319	2.469	1.892	1.508	0.949	0.293	0.137	0.088	0.016	0.038					
2006	6.846	1.914	2.500	3.237	3.351	2.146	1.328	1.108	0.277	0.146		0.008	0.015				
2007	6.561	0.746	1.935	1.730	1.710	1.233	0.636	0.534	0.137	0.082	0.004	0.023		0.002			
2008	18.407	3.291	2.689	3.568	2.550	1.912	1.271	0.758	0.436	0.200	0.068	0.030	0.032	0.005	0.005		
2009	3.670	6.757	2.715	4.221	4.487	2.045	1.147	0.803	0.615	0.256	0.081	0.092	0.005		0.002	0.006	
2010	0.957	7.954	2.777	3.931	4.038	3.054	1.019	0.518	0.735	0.378	0.167	0.138	0.018	0.014	0.004	0.015	
2011	0.477	3.763	4.715	3.812	4.001	2.977	1.828	0.374	0.435	0.256	0.118	0.056	0.149	0.008			
2012	0.400	5.463	3.970	5.075	3.391	2.092	1.341	1.001	0.504	0.225	0.210	0.046	0.029	0.043	0.017	0.002	
2013	0.944	7.727	4.770	2.383	3.144	2.867	2.217	1.493	0.685	0.297	0.121	0.184	0.107	0.004	0.006	0.003	0.005
2014	20.339	7.671	5.936	5.326	3.630	4.146	2.149	1.530	0.532	0.384	0.059	0.161	0.084	0.006	0.009	0.016	0.004
2015	0.383	6.573	6.861	5.286	4.975	2.616	2.173	0.986	0.543	0.481	0.122	0.053	0.041	0.002	0.006		0.002
2016	3.288	4.156	5.075	5.112	3.051	2.384	0.908	0.876	0.494	0.259	0.093	0.093	0.017	0.013	0.020		0.006
2017	6.268	0.849	3.209	2.733	1.805	1.395	0.882	0.395	0.254	0.171	0.043	0.090	0.018	0.006			
2018	0.147	4.355	3.451	2.099	2.116	1.371	0.945	0.355	0.207	0.159	0.089	0.045	0.022	0.005		0.004	0.003
2019	0.439	2.571	2.269	2.949	1.956	1.964	1.144	0.818	0.224	0.215	0.101	0.083	0.040	0.022	0.003	0.002	0.007
2020	0.081	0.616	1.372	1.819	1.736	1.137	0.589	0.323	0.208	0.076	0.026	0.041		0.004	0.003		

Table B.1.2. CV abundance indices

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16

2003	0.286	0.229	0.219	0.147	0.133	0.107	0.104	0.220	0.282	0.232	0.684	1.760		0.886			0.867
2004	0.240	0.257	0.128	0.154	0.169	0.192	0.227	0.211	0.349	0.384	0.397	0.922			0.970		
2005	0.684	0.243	0.226	0.248	0.208	0.158	0.216	0.206	0.281	0.377	0.579	0.718					
2006	0.216	0.185	0.285	0.200	0.224	0.153	0.116	0.198	0.247	0.306		0.867	0.830				
2007	0.275	0.233	0.400	0.353	0.232	0.201	0.169	0.216	0.269	0.411	0.867	0.956		0.806			
2008	0.245	0.163	0.234	0.161	0.126	0.097	0.161	0.138	0.186	0.338	0.282	0.419	0.462	0.850	0.794		
2009	0.621	0.157	0.137	0.176	0.126	0.123	0.150	0.136	0.214	0.234	0.334	0.340	1.282		1.353	0.477	
2010	0.537	0.087	0.163	0.151	0.183	0.140	0.164	0.225	0.271	0.247	0.181	0.374	0.548	0.752	0.831	0.440	
2011	0.483	0.342	0.221	0.268	0.178	0.193	0.176	0.236	0.194	0.265	0.344	0.610	0.364	0.628			
2012	0.620	0.128	0.297	0.211	0.138	0.181	0.164	0.165	0.223	0.220	0.197	0.258	0.429	0.564	0.652		0.771
2013	0.519	0.123	0.145	0.177	0.218	0.205	0.225	0.282	0.177	0.307	0.353	0.477	0.405	0.872	0.641	0.806	0.682
2014	0.326	0.164	0.291	0.188	0.152	0.150	0.243	0.139	0.156	0.207	0.389	0.392	0.349	0.774	0.685	0.463	0.574
2015	0.620	0.154	0.127	0.192	0.244	0.265	0.200	0.286	0.185	0.182	0.236	0.400	0.274	0.897	0.448		1.203
2016	0.495	0.198	0.122	0.122	0.095	0.087	0.131	0.153	0.201	0.205	0.287	0.315	0.420	0.669	0.564		1.066
2017	0.512	0.271	0.183	0.152	0.139	0.087	0.159	0.178	0.261	0.270	0.273	0.391	0.534	0.731			
2018	0.789	0.144	0.187	0.145	0.138	0.121	0.121	0.149	0.254	0.236	0.186	0.303	0.420	0.549		0.630	0.842
2019	0.381	0.127	0.157	0.143	0.123	0.149	0.142	0.194	0.286	0.274	0.361	0.297	0.513	0.511	0.970	0.897	0.797
2020	0.740	0.260	0.118	0.121	0.116	0.127	0.144	0.144	0.220	0.387	0.381	0.276		0.837	0.813		

Table B.1.3. Biomass indices (kilotonnes)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	0.059	0.444	1.422	3.546	6.782	7.575	9.209	4.858	3.188	1.839	0.460	0.003		0.280			0.257
2004	0.035	0.300	0.976	2.731	7.157	4.948	3.832	2.296	1.214	0.248	0.609	0.042			0.045		
2005	0.000	0.105	0.554	2.360	3.563	3.724	2.409	1.335	0.540	0.524	0.130	0.600					
2006	0.047	0.197	1.352	3.706	6.120	5.268	4.683	4.184	1.369	0.960		0.201	0.061				
2007	0.046	0.095	1.237	2.218	3.615	3.344	2.253	2.656	1.094	0.856	0.078	0.077		0.041			
2008	0.129	0.325	1.560	4.785	5.534	6.240	4.478	3.679	2.084	0.891	0.308	0.147	0.124	0.042	0.039		

2009	0.026	0.497	1.027	4.394	8.242	5.397	4.156	3.320	2.575	1.197	0.402	0.511	0.015		0.005	0.053	
2010	0.007	0.484	1.157	4.367	7.776	8.335	3.848	2.243	3.385	2.313	1.000	1.206	0.149	0.029	0.054	0.101	
2011	0.005	0.246	1.723	4.247	8.519	8.424	7.248	2.316	2.197	1.935	0.683	0.197	1.123	0.041			
2012	0.003	0.425	1.349	5.044	6.842	6.441	4.708	4.271	2.868	1.501	1.468	0.197	0.178	0.254	0.121		0.008
2013	0.011	0.638	1.868	2.284	6.150	8.462	9.972	6.377	4.333	1.699	0.700	1.331	1.302	0.020	0.050	0.042	0.086
2014	0.179	0.614	2.634	6.209	6.622	11.656	7.820	6.943	3.019	2.252	0.406	1.182	0.641	0.034	0.075	0.120	0.036
2015	0.002	0.517	3.110	6.432	10.737	7.299	7.537	4.454	3.416	2.772	0.841	0.545	0.180	0.023	0.065		0.019
2016	0.022	0.280	1.891	5.894	6.053	7.885	3.441	3.994	3.241	1.915	0.566	0.888	0.090	0.055	0.184		0.054
2017	0.031	0.074	1.561	2.963	3.482	4.457	3.505	1.721	1.842	1.152	0.250	0.976	0.097	0.085			
2018	0.001	0.249	1.779	2.333	4.184	4.289	3.516	2.017	1.285	1.218	0.623	0.212	0.098	0.039		0.074	0.023
2019	0.004	0.170	0.836	3.480	4.190	6.171	4.676	3.822	1.039	1.523	0.680	0.712	0.357	0.168	0.027	0.028	0.047
2020	0.001	0.045	0.545	1.897	3.305	3.578	2.051	1.805	1.296	0.548	0.158	0.376		0.023	0.024		

Table B.1.4. Length at age (cm)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	9.69	19.86	34.62	44.13	51.43	60.37	69.24	77.02	77.79	83.29	95.32	61.06		118.78			143.00
2004	8.97	21.32	33.97	44.88	56.38	60.30	69.51	72.16	76.45	74.73	85.78	87.83			88.94		
2005	10.97	22.58	34.77	45.23	56.19	61.91	62.55	73.38	73.35	79.01	90.27	107.44					
2006	9.38	21.82	37.77	48.09	56.62	61.73	69.12	69.81	75.40	85.66		125.00	76.00				
2007	9.20	22.55	39.17	49.46	58.43	63.90	69.13	75.48	89.59	95.22	116.00	65.07		123.00			
2008	9.16	21.76	38.19	51.09	59.41	68.28	69.74	75.52	74.14	72.32	74.63	75.35	72.40	100.00	89.00		
2009	8.66	19.63	33.16	46.77	56.59	64.32	70.88	73.50	72.55	75.67	76.48	77.81	68.97		72.98	94.96	
2010	9.59	18.17	34.72	47.50	56.93	63.86	71.93	73.91	74.50	82.06	82.01	88.38	87.68	61.46	104.00	85.23	
2011	10.24	19.25	33.54	48.18	59.34	65.19	71.74	82.77	77.55	85.42	80.34	73.91	88.09	75.38			
2012	9.39	20.46	32.48	46.80	59.04	66.77	70.76	74.60	80.30	84.34	85.31	76.60	82.78	85.38	90.46		79.00
2013	10.94	20.52	34.29	46.17	58.60	67.36	75.98	75.66	83.17	83.43	80.71	86.66	102.07	74.29	88.25	113.00	125.00
2014	10.05	20.36	35.64	49.14	56.94	65.78	71.85	77.13	81.66	81.48	84.80	87.93	85.79	83.56	85.01	86.50	93.40

2015	8.42	20.25	35.43	49.97	60.45	66.47	70.53	76.83	83.94	79.66	85.60	89.93	72.35	95.00	98.40		101.56
2016	9.30	19.09	33.58	48.49	58.81	69.78	73.30	76.15	85.26	88.16	83.07	90.66	81.80	78.15	97.36		96.42
2017	8.05	21.43	36.63	47.83	58.27	68.42	73.93	76.04	87.70	84.72	82.68	98.12	84.10	110.37			
2018	10.11	17.66	37.16	48.17	58.95	68.27	71.52	82.27	85.86	89.11	88.21	78.40	77.58	86.91		120.15	93.00
2019	9.97	19.70	33.51	49.13	59.79	68.59	74.86	77.86	78.44	85.97	86.19	93.47	92.04	92.37	96.34	109.00	96.04
2020	10.87	20.08	33.69	46.58	57.53	67.30	70.99	80.67	83.40	87.60	84.22	91.99		83.98	93.41		

Table B.1.5. Weight at age (gram)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	9	80	400	819	1332	2233	3379	4761	5346	6805	9921	2033		17080			25600
2004	7	98	391	945	1852	2350	3563	4100	5035	4538	7973	7865			7130		
2005	14	115	417	955	1884	2462	2553	4462	3959	5552	8077	14845					
2006	7	103	537	1140	1819	2457	3528	3824	4901	6596		25080	4220				
2007	7	125	659	1321	2135	2714	3540	4935	7953	10119	17480	3485		21100			
2008	7	99	575	1339	2167	3265	3524	4849	4770	4493	4541	5040	4114	9250	8520		
2009	9	74	379	1040	1838	2638	3629	4138	4190	4700	5112	5553	3082		3122	9681	
2010	7	61	418	1111	1924	2730	3772	4343	4720	6084	5958	8337	7793	2120	12154	6475	
2011	10	64	366	1123	2139	2853	3957	6170	5047	7495	5668	4111	7567	4653			
2012	7	78	335	995	2013	3064	3499	4275	5585	6622	6957	4411	6173	6953	7006		4630
2013	12	83	389	960	1962	2954	4424	4270	6290	5632	5543	7587	12149	5104	7738	16220	16300
2014	9	80	442	1168	1825	2813	3629	4533	5684	5840	6604	7543	7420	5313	7920	8237	8263
2015	4	78	453	1212	2149	2818	3480	4558	6243	5795	6695	9301	4334	10170	10132		11296
2016	7	68	371	1152	1982	3309	3793	4555	6535	7434	6152	9403	5218	4791	9660		8325
2017	5	88	486	1086	1932	3193	3975	4369	7099	6769	5913	10482	5300	13790			
2018	6	57	512	1109	1975	3125	3719	5692	6216	7726	6947	4781	4564	8512		18477	7350
2019	9	66	368	1181	2139	3130	4085	4660	4640	7008	6731	8592	8869	7996	9585	12345	7040
2020	7	75	397	1041	1901	3140	3482	5579	6216	7095	6372	9186		5165	8589		

Table B.1.6. Abundance index, standard deviation (SD) and Coefficient of variation (CV) for sum of age 2+ fish

	Index_2plus	SD_2plus	CV_2plus
2003	20.974	2.403	0.115
2004	13.357	1.682	0.126
2005	8.710	1.572	0.180
2006	14.114	2.250	0.159
2007	8.027	2.061	0.257
2008	13.522	1.587	0.117
2009	16.474	1.656	0.101
2010	16.805	2.140	0.127
2011	18.730	3.101	0.166
2012	17.947	2.661	0.148
2013	18.286	2.434	0.133
2014	23.973	4.084	0.170
2015	24.148	3.592	0.149
2016	18.404	1.537	0.084
2017	11.001	1.331	0.121
2018	10.871	1.211	0.111
2019	11.798	1.076	0.091
2020	7.335	0.567	0.077

### 3.2 - Subarea A: North of 67°N

Table B.2.1. Abundance indices (millions)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	6.238	5.254	3.268	3.763	4.521	2.700	2.319	0.863	0.489	0.220	0.042	0.001		0.016		0.010	
2004	4.997	2.837	2.201	2.396	2.602	1.463	0.722	0.359	0.181	0.046	0.050	0.006		0.006			

2005	0.026	0.665	1.042	1.988	1.478	1.268	0.746	0.157	0.107	0.068	0.016	0.038						
2006	5.896	1.802	2.156	2.623	2.946	1.554	1.026	0.941	0.171	0.107		0.008	0.015					
2007	6.561	0.446	0.911	0.853	1.071	0.789	0.465	0.394	0.114	0.075	0.004	0.022		0.002				
2008	15.128	2.463	1.822	2.795	1.883	1.419	1.145	0.580	0.348	0.161	0.052	0.011	0.021	0.005	0.005			
2009	3.454	6.642	2.251	3.570	3.716	1.584	0.868	0.712	0.466	0.204	0.076	0.072	0.004		0.002	0.006		
2010	0.424	7.412	2.353	3.268	3.385	2.397	0.784	0.383	0.733	0.317	0.151	0.137	0.018	0.002	0.004	0.015		
2011	0.308	2.322	3.471	2.498	2.866	2.095	1.445	0.292	0.315	0.213	0.116	0.055	0.130	0.008				
2012	0.388	4.299	3.218	4.485	2.784	1.537	1.042	0.930	0.411	0.200	0.209	0.046	0.029	0.043	0.017		0.002	
2013	0.486	6.382	4.101	1.706	2.666	1.887	1.575	0.890	0.578	0.297	0.110	0.184	0.107	0.004	0.006	0.003	0.005	
2014	9.259	5.696	5.448	4.026	3.034	3.521	2.016	1.388	0.465	0.364	0.059	0.159	0.083	0.006	0.009	0.016	0.004	
2015	0.381	4.298	4.733	4.154	3.727	2.068	1.818	0.902	0.506	0.397	0.119	0.053	0.040	0.002	0.006		0.002	
2016	3.233	3.944	4.433	4.522	2.610	1.995	0.746	0.735	0.413	0.203	0.067	0.093	0.017	0.005	0.020		0.006	
2017	6.268	0.768	2.891	2.407	1.563	1.151	0.715	0.308	0.200	0.147	0.043	0.090	0.018	0.006				
2018	0.135	4.070	3.197	1.916	1.879	1.049	0.748	0.323	0.183	0.128	0.089	0.045	0.022	0.005		0.004	0.003	
2019	0.148	2.234	2.114	2.470	1.508	1.460	0.839	0.490	0.148	0.129	0.075	0.069	0.035	0.021	0.003	0.002	0.007	
2020	0.066	0.571	1.149	1.520	1.367	0.861	0.499	0.210	0.129	0.042	0.025	0.040		0.004	0.003			

Table B.2.2. CV abundance indices

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	0.287	0.231	0.230	0.159	0.140	0.120	0.116	0.244	0.319	0.245	0.691	0.941		0.886			0.867
2004	0.240	0.274	0.158	0.158	0.161	0.209	0.205	0.234	0.344	0.399	0.374	0.922			0.970		
2005	0.611	0.208	0.282	0.295	0.220	0.160	0.248	0.241	0.254	0.452	0.579	0.718					
2006	0.248	0.199	0.336	0.242	0.257	0.174	0.129	0.244	0.299	0.339		0.867	0.830				
2007	0.275	0.226	0.276	0.295	0.180	0.170	0.153	0.238	0.306	0.443	0.867	0.915		0.806			
2008	0.192	0.150	0.263	0.214	0.126	0.109	0.169	0.152	0.202	0.365	0.356	0.346	0.586	0.850	0.794		
2009	0.658	0.160	0.160	0.182	0.144	0.135	0.179	0.147	0.207	0.244	0.273	0.407	0.682		0.844	0.477	
2010	0.476	0.087	0.157	0.192	0.212	0.163	0.180	0.258	0.271	0.213	0.161	0.373	0.518	0.872	0.831	0.440	

2011	0.408	0.203	0.242	0.267	0.186	0.230	0.173	0.246	0.230	0.232	0.353	0.597	0.366	0.628				
2012	0.631	0.091	0.374	0.237	0.132	0.116	0.127	0.160	0.174	0.230	0.196	0.258	0.429	0.564	0.652			0.771
2013	0.663	0.137	0.170	0.148	0.231	0.199	0.213	0.164	0.172	0.307	0.385	0.477	0.405	0.872	0.641	0.806	0.682	
2014	0.411	0.166	0.303	0.166	0.165	0.172	0.262	0.141	0.154	0.221	0.389	0.394	0.348	0.774	0.685	0.453	0.574	
2015	0.624	0.188	0.174	0.184	0.271	0.294	0.220	0.300	0.190	0.188	0.230	0.402	0.174	0.897	0.448			1.203
2016	0.499	0.197	0.126	0.134	0.104	0.093	0.130	0.158	0.237	0.202	0.196	0.315	0.420	0.527	0.564			1.066
2017	0.512	0.297	0.195	0.169	0.155	0.092	0.174	0.184	0.385	0.297	0.273	0.391	0.534	0.731				
2018	0.805	0.146	0.186	0.156	0.151	0.125	0.109	0.153	0.269	0.189	0.185	0.303	0.420	0.545		0.630	0.842	
2019	0.595	0.149	0.155	0.161	0.128	0.102	0.094	0.122	0.173	0.247	0.304	0.220	0.418	0.437	0.970	0.897	0.797	
2020	0.814	0.271	0.131	0.139	0.124	0.152	0.148	0.151	0.234	0.329	0.375	0.267		0.823	0.706			

Table B.2.3. Biomass indices (kilotonnes)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	0.059	0.441	1.309	2.981	5.981	5.664	7.962	4.249	2.840	1.442	0.453	0.002		0.280			0.257
2004	0.035	0.287	0.834	2.192	4.133	3.134	2.433	1.425	1.066	0.177	0.214	0.042		0.045			
2005	0.000	0.067	0.455	1.676	2.424	2.851	1.689	0.577	0.441	0.467	0.130	0.600					
2006	0.039	0.181	1.157	3.060	5.387	3.748	3.916	3.379	0.683	0.752		0.201	0.061				
2007	0.046	0.042	0.583	1.135	2.270	2.186	1.667	2.111	0.942	0.812	0.078	0.075		0.041			
2008	0.105	0.243	0.865	3.573	3.734	4.699	3.993	2.929	1.845	0.743	0.246	0.043	0.095	0.042	0.039		
2009	0.023	0.482	0.744	3.412	6.385	4.067	3.210	2.919	2.026	0.854	0.387	0.428	0.013		0.005	0.053	
2010	0.004	0.423	0.937	3.410	6.240	6.423	3.067	1.606	3.382	1.945	0.926	1.204	0.149	0.006	0.054	0.101	
2011	0.003	0.133	1.312	2.732	6.143	5.850	5.185	1.517	1.607	1.519	0.679	0.194	0.933	0.041			
2012	0.003	0.309	1.105	4.272	5.579	4.475	3.593	3.946	1.939	1.145	1.466	0.197	0.178	0.254	0.121		0.008
2013	0.005	0.534	1.662	1.713	5.321	5.893	5.866	3.591	3.093	1.699	0.673	1.331	1.302	0.020	0.050	0.042	0.086
2014	0.074	0.454	2.257	4.761	5.729	9.852	7.328	6.114	2.715	2.126	0.406	1.179	0.639	0.034	0.075	0.119	0.036
2015	0.002	0.337	2.282	5.151	8.598	5.904	6.803	4.167	3.300	2.186	0.831	0.544	0.176	0.023	0.065		0.019
2016	0.021	0.263	1.669	5.006	5.353	6.663	3.002	3.617	2.682	1.347	0.476	0.888	0.090	0.027	0.184		0.054

2017	0.031	0.067	1.339	2.470	2.994	3.707	2.883	1.438	1.415	0.961	0.250	0.976	0.097	0.085				
2018	0.001	0.242	1.694	2.127	3.711	3.143	2.776	1.840	1.115	1.096	0.623	0.211	0.098	0.039		0.074	0.023	
2019	0.001	0.143	0.790	2.681	2.890	4.449	3.301	2.317	0.724	0.883	0.553	0.641	0.337	0.162	0.027	0.028	0.047	
2020	0.000	0.042	0.352	1.419	2.627	2.548	1.726	1.055	0.832	0.317	0.157	0.372		0.022	0.023			

Table B.2.4. Length at age (cm)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	9.69	19.86	34.52	43.95	51.38	59.50	69.59	77.30	78.67	82.01	95.43	61.05		118.78			143.00
2004	8.98	21.36	33.66	44.33	54.01	59.14	68.71	71.61	79.70	71.48	73.81	87.83			88.94		
2005	10.96	21.73	35.51	44.17	54.42	60.43	60.45	71.41	74.38	83.28	90.27	107.44					
2006	9.29	21.60	37.56	48.50	56.86	61.62	70.83	69.49	71.54	87.71		125.00	76.00				
2007	9.20	20.57	38.99	50.03	58.86	63.84	69.77	76.79	89.42	95.97	116.00	65.20		123.00			
2008	9.21	21.88	36.48	50.50	58.37	68.79	69.74	76.98	76.30	74.10	75.26	71.91	74.37	100.00	89.00		
2009	8.31	19.55	32.28	45.69	55.67	64.07	71.00	73.52	72.96	73.82	76.91	79.39	69.00		73.00	94.96	
2010	10.54	17.93	34.24	46.78	56.25	63.65	73.25	73.24	74.52	81.68	81.95	88.49	87.77	60.09	104.00	85.23	
2011	10.39	18.66	34.05	48.06	59.55	65.11	70.59	78.70	79.45	83.87	80.60	74.00	86.43	75.38			
2012	9.35	19.99	32.61	46.25	58.88	65.58	70.11	74.24	76.02	81.24	85.38	76.60	82.78	85.38	90.46		79.00
2013	10.64	20.68	34.70	46.99	59.21	68.33	73.12	74.04	80.24	83.43	82.22	86.66	102.07	74.29	88.25	113.00	125.00
2014	9.28	20.37	34.99	49.60	57.77	65.93	71.85	76.62	82.41	81.31	84.80	88.16	85.85	83.56	85.01	86.51	93.40
2015	8.41	20.00	36.42	50.33	62.01	67.46	72.79	77.52	85.01	79.01	85.87	90.02	72.48	95.00	98.40		101.56
2016	9.26	19.08	33.66	48.12	59.56	70.51	74.95	78.42	85.00	85.34	86.21	90.66	81.80	81.11	97.36		96.42
2017	8.05	21.43	36.14	47.20	58.11	68.93	74.45	78.57	86.22	83.86	82.68	98.12	84.10	110.37			
2018	10.09	18.05	37.50	48.27	59.09	67.72	71.64	82.25	85.73	92.34	88.22	78.41	77.58	86.93		120.15	93.00
2019	9.35	19.56	33.76	48.32	58.51	68.48	74.41	78.87	80.57	86.10	89.59	95.89	92.88	92.56	96.34	109.00	96.04
2020	10.63	19.97	31.82	45.34	57.86	66.46	71.26	79.10	84.74	88.11	84.37	92.13		84.00	93.58		

Table B.2.5. Weight at age (gram)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	9	80	397	791	1322	2098	3421	4774	5559	6519	9954	2033		17080			25600
2004	7	98	378	913	1588	2154	3373	3942	5693	3751	4183	7865			7130		
2005	14	101	431	843	1649	2249	2277	3662	4139	6331	8077	14845					
2006	7	100	529	1162	1816	2410	3822	3645	4026	7059		25080	4220				
2007	7	94	629	1312	2113	2762	3581	5277	8139	10381	17480	3523		21100			
2008	7	99	470	1272	1985	3314	3489	5036	5253	4742	4719	3796	4960	9250	8520		
2009	8	73	331	953	1719	2567	3703	4102	4347	4217	5204	5874	3084		3122	9681	
2010	9	57	399	1043	1840	2685	3902	4217	4724	6131	6111	8362	7813	2635	12154	6475	
2011	10	57	377	1091	2153	2814	3589	5181	5110	7083	5714	4122	7288	4653			
2012	7	72	336	954	2004	2911	3449	4248	4712	5714	6970	4411	6173	6953	7006		4630
2013	12	84	402	1004	2003	3119	3724	4042	5371	5632	5803	7587	12149	5104	7738	16220	16300
2014	8	80	411	1179	1887	2802	3623	4402	5840	5812	6604	7585	7436	5313	7920	8240	8263
2015	4	78	480	1232	2287	2891	3756	4679	6466	5492	6759	9330	4369	10170	10132		11296
2016	7	67	375	1104	2049	3340	4025	4916	6451	6672	7009	9403	5218	5602	9660		8325
2017	5	88	462	1027	1919	3218	4031	4685	6872	6553	5913	10482	5300	13790			
2018	6	60	526	1106	1974	2996	3712	5699	6147	8461	6949	4782	4564	8517		18477	7350
2019	10	64	373	1085	1918	3053	3936	4721	4906	6788	7312	9241	9115	8042	9585	12345	7040
2020	7	75	307	931	1918	2942	3455	5006	6451	7429	6405	9235		5165	8634		

Table B.2.6. Abundance index, standard deviation (SD) and Coefficient of variation (CV) for sum of age 2+ fish

	Index_2plus	SD_2plus	CV_2plus
2003		18.212	2.288
2004		10.031	1.221
2005		6.908	1.427
2006		11.547	2.317
2007		4.700	0.828

2008		10.247		1.417		0.138
2009		13.530		1.397		0.103
2010		13.947		2.003		0.144
2011		13.505		2.247		0.166
2012		14.952		2.514		0.168
2013		14.120		1.813		0.128
2014		20.602		3.586		0.174
2015		18.528		2.987		0.161
2016		15.865		1.324		0.083
2017		9.540		1.262		0.132
2018		9.591		1.087		0.113
2019		9.368		0.877		0.094
2020		5.848		0.493		0.084

### 3.3 - Subarea B: Between 65° and 67°N

Table B.3.1. Abundance indices (millions)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	0.015	0.054	0.235	0.479	0.344	0.296	0.161	0.073	0.049	0.013	0.000	0.002		0.000		0.000	
2004	0.026	0.094	0.315	0.350	0.567	0.345	0.207	0.095	0.061	0.004	0.021	0.000			0.000		
2005	0.004	0.049	0.186	0.275	0.153	0.152	0.075	0.073	0.013	0.020	0.001	0.000					
2006	0.106	0.119	0.248	0.499	0.275	0.317	0.254	0.159	0.035	0.038		0.000	0.000				
2007	0.008	0.245	0.912	0.696	0.500	0.309	0.183	0.069	0.002	0.001	0.000	0.001		0.000			
2008	0.443	0.341	0.517	0.574	0.376	0.369	0.105	0.119	0.086	0.030	0.015	0.012	0.011	0.000	0.000		
2009	0.216	0.131	0.457	0.440	0.548	0.250	0.202	0.058	0.148	0.053	0.003	0.012	0.000		0.000	0.000	
2010	0.031	0.595	0.514	0.484	0.442	0.527	0.225	0.141	0.008	0.024	0.026	0.001	0.002	0.012	0.000	0.000	
2011	0.170	1.487	1.216	1.124	0.871	0.562	0.221	0.017	0.045	0.011	0.006	0.001	0.019	0.000			

2012	0.012	1.248	1.001	0.509	0.396	0.359	0.107	0.055	0.013	0.030	0.010	0.011	0.002	0.005	0.000		0.002
2013	0.326	1.070	0.669	0.689	0.462	0.604	0.212	0.078	0.009	0.005	0.012	0.001	0.003	0.000	0.000	0.000	0.000
2014	10.944	1.749	0.341	1.019	0.350	0.242	0.048	0.071	0.049	0.039	0.003	0.000	0.007	0.001	0.000	0.000	0.000
2015	0.002	2.208	1.299	0.966	1.030	0.454	0.286	0.087	0.028	0.076	0.025	0.002	0.007	0.000	0.002		0.000
2016	0.067	0.358	0.621	0.258	0.303	0.321	0.187	0.158	0.058	0.052	0.008	0.009	0.000	0.008	0.000		0.000
2017	0.000	0.038	0.289	0.272	0.116	0.166	0.079	0.067	0.061	0.027	0.006	0.004	0.001	0.001			
2018	0.012	0.405	0.226	0.172	0.145	0.118	0.099	0.056	0.036	0.009	0.011	0.005	0.001	0.003		0.000	0.000
2019	0.318	0.348	0.239	0.335	0.291	0.229	0.053	0.087	0.024	0.035	0.018	0.012	0.001	0.000	0.000	0.000	0.000
2020	0.015	0.019	0.139	0.240	0.302	0.119	0.091	0.048	0.033	0.012	0.002	0.003		0.000	0.002		

Table B.3.2. CV abundance indices

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	1.511	0.545	0.626	0.405	0.460	0.420	0.456	0.528	0.580	0.821		1.754					
2004	0.899	0.653	0.302	0.312	0.308	0.363	0.658	0.613	0.612	1.755	0.926						
2005	2.970	0.637	0.452	0.486	0.407	0.369	0.451	0.466	0.818	0.524	0.867						
2006	0.935	0.460	0.463	0.305	0.405	0.356	0.284	0.355	0.638	0.567							
2007	0.965	0.642	0.850	0.873	0.766	0.706	0.433	0.642	0.909	6.516		6.592					
2008	0.558	0.251	0.458	0.392	0.301	0.262	0.487	0.262	0.361	0.591	0.767	0.781	0.831				
2009	0.686	0.500	0.423	0.515	0.406	0.440	0.494	0.569	0.475	0.560	2.956	0.880	10.913		16.653		
2010	0.927	0.366	0.378	0.347	0.306	0.336	0.358	0.443	0.885	0.599	0.582	6.463	1.592	0.823			
2011	0.928	0.680	0.556	0.831	0.625	0.572	0.535	1.064	0.621	1.927	1.273	5.298	0.903				
2012	2.247	0.475	0.315	0.235	0.395	0.397	0.360	0.397	0.620	0.787	0.757	0.670	0.778	0.835		0.771	
2013	0.811	0.531	0.260	0.371	0.403	0.504	0.417	0.487	0.069	0.460	0.686	0.776	0.395				
2014	0.455	0.405	0.307	0.548	0.467	0.445	0.479	0.683	0.493	0.702	0.520		0.979	0.803			
2015	4.653	0.354	0.318	0.523	0.522	0.416	0.426	0.517	0.717	0.701	0.675	1.683	1.217		0.791		
2016	1.410	0.538	0.315	0.434	0.344	0.289	0.387	0.376	0.584	0.593	0.700	0.403		0.944			
2017		0.655	0.293	0.339	0.313	0.304	0.338	0.305	0.511	0.369	0.678	0.198	0.697	0.859			

2018	1.958	0.624	0.467	0.323	0.321	0.297	0.271	0.263	0.362	0.441	0.468	0.620	0.806	0.657				
2019	0.475	0.482	0.358	0.354	0.471	0.382	0.405	0.359	0.459	0.430	0.662	0.601	1.141					
2020	1.198	1.020	0.374	0.392	0.465	0.594	0.551	0.503	0.633	0.548	1.342	1.212		10.862	1.097			

Table B.3.3. Biomass indices (kilotonnes)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	0.000	0.004	0.090	0.324	0.371	0.537	0.406	0.212	0.146	0.031	0.000	0.003		0.000			0.000
2004	0.000	0.009	0.140	0.273	0.995	0.635	0.561	0.283	0.156	0.011	0.394	0.000			0.000		
2005	0.000	0.004	0.060	0.168	0.197	0.303	0.191	0.214	0.038	0.057	0.014	0.000					
2006	0.001	0.016	0.115	0.444	0.349	0.539	0.593	0.690	0.205	0.207		0.000	0.000				
2007	0.000	0.043	0.441	0.596	0.772	0.702	0.570	0.304	0.012	0.003	0.000	0.002		0.000			
2008	0.005	0.029	0.267	0.765	0.764	1.071	0.397	0.398	0.237	0.125	0.061	0.044	0.029	0.000	0.000		
2009	0.003	0.011	0.170	0.533	0.960	0.626	0.561	0.149	0.546	0.345	0.010	0.025	0.001		0.000	0.000	
2010	0.000	0.041	0.235	0.479	0.870	1.313	0.628	0.657	0.021	0.067	0.125	0.002	0.005	0.023	0.000	0.000	
2011	0.001	0.116	0.366	1.045	1.418	1.285	0.583	0.044	0.420	0.026	0.019	0.004	0.190	0.000			
2012	0.000	0.123	0.306	0.376	0.619	1.033	0.304	0.183	0.053	0.372	0.038	0.056	0.009	0.021	0.000		0.008
2013	0.004	0.057	0.181	0.513	0.672	1.361	0.465	0.214	0.043	0.021	0.031	0.005	0.014	0.000	0.000	0.000	
2014	0.104	0.146	0.187	0.825	0.341	0.532	0.133	0.382	0.228	0.150	0.013	0.000	0.024	0.003	0.000	0.000	
2015	0.000	0.171	0.344	1.004	1.628	1.025	0.628	0.287	0.086	0.570	0.091	0.010	0.029	0.000	0.013		0.000
2016	0.001	0.029	0.184	0.169	0.426	0.759	0.499	0.470	0.269	0.520	0.038	0.052	0.000	0.028	0.000		0.000
2017	0.000	0.003	0.130	0.280	0.213	0.393	0.283	0.228	0.453	0.208	0.024	0.017	0.003	0.014			
2018	0.000	0.014	0.073	0.172	0.226	0.287	0.295	0.238	0.173	0.046	0.050	0.018	0.008	0.019		0.000	0.000
2019	0.003	0.028	0.063	0.373	0.598	0.565	0.158	0.309	0.074	0.152	0.082	0.065	0.005	0.000	0.000	0.000	
2020	0.000	0.002	0.055	0.254	0.524	0.289	0.294	0.219	0.153	0.052	0.006	0.015		0.000	0.009		

Table B.3.4. Length at age (cm)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
--	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	--------	--------	--------	--------	--------	--------	--------

2003	10.00	19.86	34.38	40.64	47.42	56.53	63.37	66.45	68.28	64.26		61.05						
2004	9.00	22.14	35.01	43.03	54.24	56.32	61.95	65.45	63.70	72.18	120.83							
2005	10.63	20.97	31.37	39.79	50.01	58.34	62.15	64.65	63.88	64.78	108.00							
2006	11.67	24.94	35.74	44.65	49.84	55.38	61.70	69.45	81.23	80.26								
2007	9.97	25.38	34.29	43.01	52.39	60.32	65.77	74.05	91.00	60.77		58.38						
2008	11.68	21.15	37.10	50.48	57.89	65.43	69.66	65.79	65.10	70.71	72.87	70.87	69.97					
2009	11.49	21.64	32.15	47.96	54.65	61.33	66.52	63.82	70.60	80.70	65.61	59.17	67.40		70.00			
2010	12.00	19.44	35.12	44.92	56.90	61.39	64.88	75.02	61.25	66.95	81.66	60.93	60.01	62.00				
2011	9.83	20.16	30.42	43.37	55.75	61.78	64.03	65.15	87.65	60.71	69.19	62.05	102.00					
2012	10.65	22.32	31.60	42.93	54.93	67.05	67.09	68.26	73.69	93.93	74.17	78.31	73.00	74.94			79.00	
2013	11.44	17.74	30.70	42.83	52.60	61.46	62.94	65.93	76.37	76.09	63.67	69.00	74.70					
2014	10.73	20.46	38.23	42.69	46.62	58.93	67.47	77.68	75.01	74.64	81.28		71.69	75.00				
2015	10.88	20.51	29.80	46.53	54.35	60.63	60.50	70.40	71.33	85.50	74.51	76.85	75.82		95.00			
2016	10.35	20.58	31.65	39.51	52.06	60.54	64.65	66.41	77.25	97.35	75.60	83.76		73.00				
2017		22.16	36.00	47.14	56.71	62.37	70.89	69.08	87.12	88.21	76.19	76.05	78.00	99.00				
2018	10.66	15.21	32.88	45.33	53.87	61.84	66.36	73.71	77.41	78.04	78.23	75.19	98.00	80.87				
2019	10.62	20.29	29.52	47.40	58.30	62.36	65.12	71.49	68.09	73.51	73.95	86.65	76.97					
2020	11.94	22.83	34.55	46.89	55.10	62.06	66.73	75.66	77.61	77.43	78.48	82.85		81.80	82.97			

Table B.3.5. Weight at age (gram)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	10	69	383	667	1070	1831	2592	2876	3172	2627		2033					
2004	6	107	440	772	1737	1823	2484	2861	2519	3396	19004						
2005	14	85	329	617	1278	1964	2527	2941	2671	2873	12140						
2006	13	139	457	887	1257	1693	2336	4269	5914	5353							
2007	7	160	422	806	1518	2250	3138	4143	6804	2281		1485					
2008	11	85	526	1322	2058	2916	3785	3412	2749	4571	3955	3716	2737				

2009	14	86	354	1152	1734	2474	2848	2605	3606	6205	3050	2022	3084		3122			
2010	14	69	445	978	1973	2469	2793	4551	2362	2856	4925	2825	2285	1915				
2011	9	75	285	818	1683	2361	2681	2839	9183	2343	3580	2440	10116					
2012	9	100	311	736	1555	2912	2804	3312	4149	9801	4212	4856	3745	4318				4630
2013	12	52	272	760	1460	2193	2240	2739	4584	4326	2554	3425	4262					
2014	10	86	534	739	973	2161	2876	4915	4475	4035	4659		3635	3340				
2015	19	77	263	1000	1554	2278	2241	3392	3418	7843	3516	4276	4335		7785			
2016	9	84	294	661	1411	2376	2678	3003	4680	10566	4579	5951		3425				
2017		92	444	1020	1844	2366	3567	3390	6677	7467	4049	4314	3935	10610				
2018	5	40	327	1010	1585	2468	2987	4217	4882	4988	4720	3535	8695	6416				
2019	11	77	267	1100	2005	2497	2973	3577	3220	4252	4282	6269	4317					
2020	7	88	393	1056	1735	2463	3166	4680	4637	4613	3372	6073		5165	5434			

Table B.3.6. Abundance index, standard deviation (SD) and Coefficient of variation (CV) for sum of age 2+ fish

	Index_2plus	SD_2plus	CV_2plus
2003		1.650	0.655
2004		1.965	0.651
2005		0.949	0.377
2006		1.826	0.508
2007		2.673	2.045
2008		2.215	0.620
2009		2.171	0.888
2010		2.406	0.727
2011		4.092	2.531
2012		2.500	0.574
2013		2.745	0.718
2014		2.170	0.680

2015		4.262		1.573		0.369
2016		1.982		0.561		0.283
2017		1.088		0.270		0.248
2018		0.883		0.233		0.263
2019		1.323		0.401		0.303
2020		0.990		0.415		0.419

### 3.4 - Subarea C: Between 62°N and 65°N

Table B.4.1. Abundance indices (millions)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	0.000	0.010	0.058	0.260	0.284	0.429	0.254	0.077	0.055	0.042	0.001	0.001		0.000		0.000	
2004	0.054	0.065	0.074	0.235	0.781	0.355	0.161	0.110	0.004	0.009	0.000	0.000			0.000		
2005	0.000	0.191	0.107	0.230	0.285	0.122	0.135	0.069	0.022	0.000	0.001	0.000					
2006	0.844	0.007	0.115	0.116	0.157	0.324	0.073	0.024	0.071	0.000		0.000	0.000				
2007	0.008	0.073	0.146	0.198	0.149	0.155	0.031	0.083	0.025	0.006	0.000	0.000		0.000			
2008	2.836	0.487	0.349	0.199	0.291	0.125	0.021	0.058	0.001	0.009	0.000	0.006	0.000	0.000	0.000		
2009	0.000	0.083	0.226	0.289	0.340	0.215	0.083	0.049	0.011	0.005	0.002	0.007	0.000		0.000	0.000	
2010	0.503	0.451	0.092	0.306	0.286	0.234	0.035	0.006	0.007	0.051	0.010	0.000	0.002	0.000	0.000	0.000	
2011	0.000	0.047	0.120	0.241	0.286	0.326	0.170	0.083	0.075	0.031	0.004	0.000	0.000	0.000			
2012	0.000	0.083	0.308	0.374	0.261	0.229	0.231	0.077	0.104	0.004	0.009	0.011	0.002	0.005	0.000	0.002	
2013	0.131	0.318	0.183	0.120	0.087	0.408	0.470	0.545	0.116	0.005	0.001	0.001	0.003	0.000	0.000	0.000	
2014	0.155	0.354	0.217	0.302	0.258	0.411	0.126	0.110	0.038	0.060	0.003	0.002	0.007	0.001	0.000	0.000	
2015	0.000	0.086	0.839	0.184	0.250	0.098	0.095	0.005	0.024	0.027	0.022	0.002	0.006	0.000	0.002	0.000	
2016	0.012	0.147	0.122	0.393	0.197	0.196	0.074	0.070	0.074	0.021	0.035	0.009	0.000	0.000	0.000	0.000	
2017	0.000	0.108	0.163	0.107	0.143	0.143	0.124	0.068	0.007	0.004	0.006	0.004	0.001	0.001			
2018	0.000	0.120	0.082	0.068	0.113	0.259	0.193	0.042	0.049	0.039	0.011	0.005	0.001	0.003	0.000	0.000	

2019	0.028	0.012	0.084	0.183	0.177	0.296	0.256	0.271	0.064	0.061	0.015	0.012	0.006	0.001	0.000	0.000	0.000
2020	0.000	0.025	0.140	0.133	0.100	0.198	0.036	0.077	0.055	0.032	0.001	0.002		0.000	0.001		

Table B.4.2. CV abundance indices

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003		0.712	0.779	0.483	0.829	0.529	0.715	0.860	0.934	0.813	5.884	0.830					
2004	1.250	0.998	0.463	0.415	0.407	0.389	0.429	0.549	0.936	0.790	7.491						
2005		0.636	0.609	0.530	0.448	0.570	0.478	0.658	0.798		0.867						
2006	0.627	0.411	0.387	0.641	0.394	0.548	0.571	0.615	0.645								
2007	0.965	0.654	0.486	0.403	0.551	0.475	0.342	0.674	0.673	0.885							
2008	0.913	0.852	0.454	0.475	0.491	0.435	0.922	0.733	4.512	0.893	8.656	0.928					
2009		0.465	0.289	0.611	0.588	0.786	0.926	0.803	1.817	2.604	5.625	0.996	13.536				
2010	0.971	0.530	0.216	0.520	0.592	0.556	0.409	0.421	0.573	0.837	0.342		0.742				
2011		0.185	0.575	0.459	0.452	0.453	0.531	0.474	0.553	0.893	0.509						
2012		0.287	0.636	0.361	0.496	0.694	0.577	0.534	0.818	0.461	0.350	0.670	0.778	0.835		0.771	
2013	0.859	0.867	0.418	0.491	0.574	0.843	0.713	0.692	0.556	0.460	0.772	0.776	0.395				
2014	1.145	0.840	0.464	0.571	0.698	0.580	0.502	0.698	0.613	0.619	0.520	5.910	1.159	0.803		22.361	
2015		0.849	0.559	0.447	0.479	0.555	0.656	0.909	0.652	0.587	0.554	0.774	0.102		0.791		
2016	0.791	0.414	0.549	0.406	0.429	0.346	0.436	0.547	0.483	0.614	0.701	0.403					
2017		0.719	0.416	0.417	0.493	0.378	0.457	0.406	0.222	0.345	0.678	0.198	0.697	0.859			
2018		0.418	0.629	0.384	0.454	0.370	0.355	0.259	0.422	0.716	0.469	0.614	0.806	0.653			
2019	0.958	0.850	0.371	0.406	0.659	0.733	0.548	0.563	0.905	0.692	1.559	1.565	2.276	5.787			
2020		0.956	0.413	0.422	0.413	0.245	0.471	0.391	0.555	0.771	0.834	0.799			0.836		

Table B.4.3. Biomass indices (kilotonnes)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003	0.000	0.000	0.028	0.299	0.469	1.401	0.872	0.424	0.274	0.380	0.006	0.002		0.000		0.000	

2004	0.000	0.004	0.019	0.308	2.092	1.227	0.853	0.595	0.009	0.075	0.001	0.000				0.000		
2005	0.000	0.034	0.043	0.525	0.966	0.611	0.538	0.557	0.070	0.000	0.014	0.000						
2006	0.006	0.001	0.085	0.202	0.408	1.043	0.235	0.138	0.481	0.000		0.000	0.000					
2007	0.000	0.012	0.218	0.495	0.581	0.490	0.116	0.273	0.164	0.041	0.000	0.000		0.000				
2008	0.019	0.053	0.427	0.448	1.036	0.471	0.088	0.352	0.002	0.022	0.000	0.059	0.000	0.000	0.000			
2009	0.000	0.009	0.158	0.480	0.969	0.710	0.397	0.299	0.029	0.013	0.005	0.058	0.001		0.000	0.000		
2010	0.002	0.048	0.016	0.523	0.720	0.736	0.210	0.020	0.018	0.337	0.052	0.000	0.004	0.000	0.000	0.000		
2011	0.000	0.003	0.061	0.488	0.973	1.296	1.484	0.808	0.170	0.391	0.014	0.000	0.000	0.000				
2012	0.000	0.008	0.121	0.555	0.687	0.995	0.905	0.339	0.976	0.016	0.037	0.056	0.009	0.021	0.000		0.008	
2013	0.002	0.050	0.064	0.125	0.213	1.277	3.744	2.639	1.283	0.021	0.004	0.005	0.014	0.000	0.000	0.000	0.000	
2014	0.002	0.024	0.202	0.632	0.564	1.326	0.489	0.593	0.150	0.276	0.013	0.004	0.026	0.003	0.000	0.000	0.000	
2015	0.000	0.010	0.485	0.283	0.533	0.372	0.192	0.021	0.103	0.110	0.084	0.009	0.025	0.000	0.013		0.000	
2016	0.000	0.012	0.064	0.754	0.331	0.811	0.272	0.253	0.528	0.190	0.129	0.052	0.000	0.000	0.000		0.000	
2017	0.000	0.009	0.128	0.248	0.301	0.522	0.446	0.211	0.026	0.017	0.024	0.017	0.003	0.014				
2018	0.000	0.007	0.026	0.067	0.282	1.054	0.773	0.182	0.297	0.168	0.050	0.018	0.008	0.019		0.000	0.000	
2019	0.001	0.001	0.016	0.439	0.726	1.185	1.232	1.298	0.295	0.523	0.071	0.072	0.024	0.006	0.000	0.000	0.000	
2020	0.000	0.002	0.156	0.267	0.204	0.841	0.144	0.580	0.355	0.233	0.005	0.011		0.000	0.008			

Table B.4.4. Length at age (cm)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003		17.31	33.13	44.67	52.18	66.76	68.29	75.08	77.81	93.01	84.42	61.04					
2004	9.16	16.91	28.55	48.68	63.18	66.87	77.57	79.22	62.80	88.52	63.44						
2005		26.78	33.20	59.22	68.01	74.51	72.93	81.79	66.42		108.00						
2006	9.98	21.96	41.84	53.82	60.92	66.50	68.23	66.22	86.82								
2007	9.97	23.65	50.34	60.55	68.31	69.03	67.63	70.49	91.31	89.00							
2008	8.33	21.34	48.25	60.00	68.10	70.79	71.94	78.57	61.38	59.05	62.00	87.89					
2009		22.22	38.04	51.59	61.81	70.70	75.86	79.22	62.77	62.25	66.95	91.73	67.00				

2010	8.02	20.70	27.14	50.94	59.27	63.95	78.94	66.91	61.27	81.12	79.67		60.00					
2011		20.51	34.19	56.06	68.10	70.66	89.96	95.59	63.83	106.00	69.52							
2012		21.72	33.95	51.94	63.37	70.99	72.63	74.78	90.09	71.36	74.33	78.31	73.00	74.94			79.00	
2013	11.00	24.25	32.75	44.82	55.34	71.04	88.43	79.79	95.79	76.09	63.00	69.00	74.70					
2014	11.70	21.12	43.63	58.48	59.96	67.83	73.87	79.82	74.50	77.93	81.28	61.71	71.66	75.00		70.00		
2015		22.25	38.86	53.28	57.32	70.80	61.01	71.05	75.34	76.42	74.87	77.00	76.15		95.00			
2016	10.00	21.14	36.26	56.79	54.97	72.89	71.14	69.90	84.50	97.40	74.29	83.76						
2017		21.85	40.91	58.32	61.16	68.15	70.64	66.42	73.92	81.74	76.19	76.05	78.00	99.00				
2018		19.52	32.19	45.60	63.57	74.71	72.80	74.12	82.67	75.60	78.26	75.22	98.00	80.88				
2019	11.83	19.97	28.09	59.05	70.49	71.02	78.73	77.01	76.47	90.06	77.19	92.56	76.89	81.69				
2020		21.00	47.03	54.73	59.39	73.19	70.63	86.06	83.71	86.60	79.00	83.00			83.00			

Table B.4.5. Weight at age (gram)

	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12	Age 13	Age 14	Age 15	Age 16
2003		40	365	1101	1436	3168	3272	5072	5062	8818	6215	2033					
2004	7	48	254	1247	2604	3510	5123	5797	2414	7569	2473						
2005		188	402	2201	3380	4439	3947	6824	2778		12140						
2006	8	81	718	1798	2558	3228	3171	4489	7765								
2007	7	169	1478	2435	3582	3182	3570	3952	7115	7500							
2008	7	99	1169	2282	3468	3720	4226	5348	2369	2334	2475	9590					
2009		99	663	1468	2691	3455	4183	5423	2703	2687	2746	7945	2870				
2010	4	95	175	1551	2360	3011	5553	3154	2369	5343	5091		2285				
2011		65	423	1909	3336	3730	8552	9298	2319	12482	3640						
2012		93	424	1487	2530	3851	3712	4126	7459	3729	4235	4856	3745	4318		4630	
2013	13	129	330	923	1976	3183	7125	4835	10222	4326	2400	3425	4262				
2014	14	74	905	2078	2133	3194	4110	5267	3894	4804	4659	2441	3634	3340		4400	
2015		97	629	1491	2113	3576	2284	3228	4368	4328	3560	4310	4395		7785		

2016	9	83	452	1899	1624	4040	3585	3558	6659	9035	3961	5951							
2017		90	758	2215	2137	3506	3509	3183	3885	4583	4049	4314	3935	10610					
2018		61	305	953	2558	4136	3965	4332	5991	4510	4724	3539	8695	6419					
2019	23	67	195	2364	4053	3666	4952	4771	4592	8250	4122	7605	4331	5340					
2020		78	1075	1866	2050	4258	3767	7391	6500	6694	3386	6140			5434				

Table B.4.6. Abundance index, standard deviation (SD) and Coefficient of variation (CV) for sum of age 2+ fish

	Index_2plus	SD_2plus	CV_2plus
2003		1.461	0.803
2004		1.728	0.590
2005		0.970	0.358
2006		0.880	0.271
2007		0.792	0.254
2008		1.060	0.362
2009		1.227	0.690
2010		1.029	0.429
2011		1.336	0.487
2012		1.616	0.622
2013		1.940	1.033
2014		1.533	0.739
2015		1.553	0.665
2016		1.191	0.285
2017		0.771	0.219
2018		0.864	0.244
2019		1.426	0.700
2020		0.776	0.135

## 12 - Appendix C

R-script to produce the time series tables and figures from Stocks bootstrap output files. This is a version for acoustic time series.

```
# espen.johnsen@hi.no, September 2019
#####
# Coastal cod by harald@hi.no and johanna.fall@hi.no sept 2020
# Oppdatert 01.11.2021 med 2020 indeks. Hele serien kjørt ut på nytt til
# //delphi/Felles/421-Bunnfisk/Kysttorsk/Acoustics/N67/, Tot, S65, osv.
# Oppdatert 27.01.2021 med SSB estimat
# Oppdatert 12.03.2021 med linje 51-55 for ? legge til nullverdier i beregningene.
# All non-coastal cod (otolithtype > 2) are stripped from the dataset, so that all output is for coastal cod only
rm(list=ls())
library(data.table)
library(Hmisc)
library(Rstox)
library(xlsx)
## Coastal cod
#path1 <- "C:Users//harald//workspace//stox//project//Varanger Stad cod acoustic index in autumn "
#path1 <- "C:Users//harald//workspace//stox//project//Varanger Stad coastal cod swept area "
path1 <- "//delphi/Felles/421-Bunnfisk/Kysttorsk/Akustikk stox/Varanger Stad cod acoustic index in autumn "
#path2 <- "D://Documents//OwnCloud//Softwarespesifikk//R//Projects//Kysttorsk//Swept area//Tot//"
#path2 <- "D://Documents//OwnCloud//Softwarespesifikk//R//Projects//Kysttorsk//Acoustics//Tot//"
# path2 <- "//delphi/Felles/421-Bunnfisk/Kysttorsk/Acoustics/N67//"
# path2 <- "//delphi/Felles/421-Bunnfisk/Kysttorsk/Acoustics/65-67//"
# path2 <- "//delphi/Felles/421-Bunnfisk/Kysttorsk/Acoustics/S65//"
#for SSB calcs - management plan
#Correction of indices - adding zeros at age for iterations with zeroes (were NA)
# path2 <- "//delphi/Felles/421-Bunnfisk/Kysttorsk/Acoustics/N-WIPaddzeroes//"
# path2 <- "//delphi/Felles/421-Bunnfisk/Kysttorsk/Acoustics/Tot-WIPaddzeroes//"
```

```
# path2 <- "//delphi/Felles/421-Bunnfisk/Kysttorsk/Acoustics/65-67-WIPaddzeroes//"
path2 <- "//delphi/Felles/421-Bunnfisk/Kysttorsk/Acoustics/S65-WIPaddzeroes//"
out <- list()
for(y in 1:26){
  year <- 1994+y
  print(year)
  load(paste0(path1,year," N67\\output\\r\\data\\bootstrapImpute.RData"))
  bootstrapVariable.out <- eval(parse(text="bootstrapImpute"))
  DT <- rbindlist(bootstrapVariable.out[[2]],idcol=TRUE)
  DT <- DT[DT$otolithtype <= 2] # Only coastal cod (otolithtype 1 and 2) included
  ## Some SuperIND may miss weight. Generate weight from a length weight relation
  id1 <- DT$weight > 0
  lm.wl <- lm(log(DT$weight[id1]) ~ log(DT$length[id1]))
  a <- exp(as.numeric(lm.wl$coeff[1]))
  b <- as.numeric(lm.wl$coeff[2])
  DT$weight[is.na(DT$weight) | DT$weight == 0] <- a*(DT$length[is.na(DT$weight) | DT$weight == 0] )^b
  #plot(DT$weight ~ DT$length)
  #Create zero values for ages not observed in an iteration
  setkey(DT, Stratum, age, .id)
  DT <- DT[CJ(Stratum, age, .id, unique=TRUE)]
  DT$Abundance[is.na(DT$Abundance)] <- 0
  DT$weight[is.na(DT$weight)] <- 0 #to include zeroes in biomass calc
  ## Output by age and by sum over age 2 to maxage
  strata <- unique(DT[,c("Stratum", "includeintotal")])
  # strata.incl <- strata[strata$includeintotal=="TRUE",]
  #manually remove strata so we only have area N67 left
  # strata.incl <- strata[!Stratum %in% c("Helgeland indre", "Helgeland ytre",
  # "Stad Halten indre", "Stad Halten ytre",
  # "Vestfjord ost"),]
```

```
#manually remove strata so we only have area 6567 left

strata.incl <- strata[!Stratum %in% c("Alta","Andfjord Vaagsfjord",
 "Fugloybanken","Hjelmsoy indre",
 "Hjelmsoy Loppa","Kvenangen",
 "Laksefjord","Malangen","Osthavet",
 "Porsangen","Soroya indre",
 "Stad Halten indre","Stad Halten ytre",
 "Tana","Ullsfjord Lyngen","Varangerfjord",
 "Vesteralen","Vestfjord indre","Vestfjord ost",
 "Vestfjord vest"),]

#manually remove strata so we only have area S65 left

strata.incl <- strata[!Stratum %in% c("Alta","Andfjord Vaagsfjord",
 "Fugloybanken","Helgeland indre",
 "Helgeland ytre","Hjelmsoy indre",
 "Hjelmsoy Loppa","Kvenangen",
 "Laksefjord","Malangen","Osthavet",
 "Porsangen","Soroya indre",
 "Tana","Ullsfjord Lyngen","Varangerfjord",
 "Vesteralen","Vestfjord indre","Vestfjord ost",
 "Vestfjord vest"),]

#manually remove strata so we have Total area left

# strata.incl <- strata[!Stratum %in% c("Vestfjord ost"),]

byGrp1 <- c(".id")

byGrp2 <- c("age", ".id")

tmp1 <- DT[lage %in% c(0,1) & Stratum %in% strata.incl$Stratum,.(Ab.Sum =sum(Abundance)),by=byGrp1]

tmp2 <- DT[Stratum %in% strata.incl$Stratum,.(Ab.Sum =sum(Abundance)),by=byGrp2]

tmp3 <- DT[Stratum %in% strata.incl$Stratum,.(Tonnes =sum(Abundance * (weight))/1000000),by=byGrp2] #

Tonnes

#For SSB calculation
```

```
tmp3b <- DT[Stratum %in% strata.incl$Stratum & stage %in% c(2,3,4),.(Tonnes =sum(Abundance *  
(weight))/1000000),by=byGrp2] # Tonnes  
  
tmp4 <- DT[Stratum %in% strata.incl$Stratum,.(ind.weight = weighted.mean(weight,Abundance)),by=byGrp2]  
  
tmp5 <- DT[Stratum %in% strata.incl$Stratum,.(ind.length = weighted.mean(length,Abundance)),by=byGrp2]  
  
#Maturity at age  
  
tmp6 <- DT[Stratum %in% strata.incl$Stratum,  
.(maturity = length(stage[stage %in% c(2,3,4)])/length(stage[stage %in% c(1,2,3,4)])),by=byGrp2]  
  
tsnTotWithout01 <- tmp1[, .("Ab.Sum.5%" = quantile(Ab.Sum, probs = .05),  
"Ab.Sum.50%" = quantile(Ab.Sum, probs = .50),  
"Ab.Sum.95%" = quantile(Ab.Sum, probs = .95),  
Ab.Sum.mean = mean(Ab.Sum),  
Ab.Sum.sd = sd(Ab.Sum),  
Ab.Sum.cv = sd(Ab.Sum)/mean(Ab.Sum))]  
  
tsnByAge <- tmp2[, .("Ab.Sum.5%" = quantile(Ab.Sum, probs = .05),  
"Ab.Sum.50%" = quantile(Ab.Sum, probs = .50),  
"Ab.Sum.95%" = quantile(Ab.Sum, probs = .95),  
Ab.Sum.mean = mean(Ab.Sum),  
Ab.Sum.sd = sd(Ab.Sum),  
Ab.Sum.cv = sd(Ab.Sum)/mean(Ab.Sum))  
, by = c("age")]  
  
tsbByAge <- tmp3[, .("Ton.5%" = quantile(Tonnes, probs = .05),  
"Ton.50%" = quantile(Tonnes, probs = .50),  
"Ton.95%" = quantile(Tonnes, probs = .95),  
Ton.mean = mean(Tonnes),  
Ton.sd = sd(Tonnes),  
Ton.cv = sd(Tonnes)/mean(Tonnes))  
, by = c("age")]  
  
ssbByAge <- tmp3b[, .("Ton.5%" = quantile(Tonnes, probs = .05),  
"Ton.50%" = quantile(Tonnes, probs = .50),
```

```
"Ton.95%" = quantile(Tonnes, probs = .95),  
Ton.mean = mean(Tonnes),  
Ton.sd = sd(Tonnes),  
Ton.cv = sd(Tonnes)/mean(Tonnes)  
, by = c("age")]  
  
ind.weight.by.age <- tmp4[, .("Ind.weight.age.5%" = quantile(ind.weight, probs = .05,na.rm=T),  
"Ind.weight.age.50%" = quantile(ind.weight, probs = .50,na.rm=T),  
"ind.weight.age.95%" = quantile(ind.weight, probs = .95,na.rm=T),  
ind.weight.age.mean = mean(ind.weight,na.rm=T),  
ind.weight.age = sd(ind.weight,na.rm=T),  
ind.weight.age.cv = sd(ind.weight,na.rm=T)/mean(ind.weight,na.rm=T))  
, by = c("age")]  
  
ind.length.by.age <- tmp5[, .("Ind.length.age.5%" = quantile(ind.length, probs = .05,na.rm=T),  
"Ind.length.age.50%" = quantile(ind.length, probs = .50,na.rm=T),  
"ind.length.age.95%" = quantile(ind.length, probs = .95,na.rm=T),  
ind.length.age.mean = mean(ind.length,na.rm=T),  
ind.length.age = sd(ind.length,na.rm=T),  
ind.length.age.cv = sd(ind.length,na.rm=T)/mean(ind.length,na.rm=T))  
, by = c("age")]  
  
maturity.by.age <- tmp6[, .("Maturity.age.5%" = quantile(maturity, probs = .05,na.rm=T),  
"Maturity.age.50%" = quantile(maturity, probs = .50,na.rm=T),  
"Maturity.age.95%" = quantile(maturity, probs = .95,na.rm=T),  
maturity.age.mean = mean(maturity,na.rm=T),  
maturity.age.sd = sd(maturity,na.rm=T),  
maturity.age.cv = sd(maturity,na.rm=T)/mean(maturity,na.rm=T))  
, by = c("age")]  
  
## Output by stratum---  
byGrp3 <- c("Stratum", ".id")  
tmp2 <- DT[ ,.(Ab.Sum =sum(Abundance)),by=byGrp3]
```

```
tmp3 <- DT[ ,.(Tonnes =sum(Abundance * (weight))/1000000),by=byGrp3] # Tonnes
tsnByStratum <- tmp2[, .("Ab.Sum.5%" = quantile(Ab.Sum, probs = .05),
"Ab.Sum.50%" = quantile(Ab.Sum, probs = .50),
"Ab.Sum.95%" = quantile(Ab.Sum, probs = .95),
Ab.Sum.mean = mean(Ab.Sum),
Ab.Sum.sd = sd(Ab.Sum),
Ab.Sum.cv = sd(Ab.Sum)/mean(Ab.Sum))
, by = c("Stratum")]

tsbByStratum <- tmp3[, .("Ton.5%" = quantile(Tonnes, probs = .05),
"Ton.50%" = quantile(Tonnes, probs = .50),
"Ton.95%" = quantile(Tonnes, probs = .95),
Ton.mean = mean(Tonnes),
Ton.sd = sd(Tonnes),
Ton.cv = sd(Tonnes)/mean(Tonnes))
, by = c("Stratum")]

#
out[[y]] <- list(year=year,
a = a,
b = b,
tsnByAge = tsnByAge,
tsnTotWithout01 = tsnTotWithout01,
ind.weight.by.age = ind.weight.by.age,
ind.length.by.age = ind.length.by.age,
maturity.by.age = maturity.by.age,
tsbByAge = tsbByAge,
ssbByAge = ssbByAge,
tsbByStratum = tsbByStratum,
tsnByStratum = tsnByStratum)
}
```

```
# If you want to save output

save(out, file = (paste0(path2,"ProduceAllFiguresCoastalCod_12-03-filtfromN.Rdata")))

out.acu <- out # I rename the object, as the "out" is in conflict with results of other codes

## Figures to be made

# Figure 1: Number by age in time series: Age1-Age17. Age 17 is the oldest observed coastal cod.

## Tables to be made

# Table 1: Abundance indices (mean bootstrap)

# Table 2: CV of abundance indices

# Table 3: Biomass indices (mean bootstrap)

# Table 4: Mean length by age (cm)

# Table 5: Mean weight by age (g)

# Table 6: Abundance and CV of abundance for sum over ages 2+

## Number by age in time series For Table 1 and Figure 1

mat.num <- matrix(NA,ncol=18,nrow=length(1995:2020))

colnames(mat.num) <- 0:17

rownames(mat.num) <- 1995:2020

for(y in 1:26){

  tmp <- (as.data.frame(out.acu[[y]]$tsnByAge)[,c("age","Ab.Sum.mean")])

  for(a in 1:18){

    age <- a-1

    if(!any(tmp$age[!is.na(tmp$age)] == age)) {

      next

    }

    mat.num[y,a] <- tmp$`Ab.Sum.mean`[tmp$age == age & !is.na(tmp$age)]

  }

}

## Biomass by age in time series. For Table 3

mat.biom <- matrix(NA,ncol=18,nrow=length(1995:2020))

colnames(mat.biom) <- 0:17
```

```
rownames(mat.biom) <- 1995:2020

for(y in 1:26){

tmp <- (as.data.frame(out.acu[[y]]$tsbByAge)[,c("age","Ton.mean")])

for(a in 1:18){

age <- a-1

if(!any(tmp$age[!is.na(tmp$age)] == age)) {

next

}

mat.biom[y,a] <- tmp$`Ton.mean`[tmp$age == age & !is.na(tmp$age)]

}

}

#SSB by age (management plan)

mat.ssb <- matrix(NA,ncol=18,nrow=length(1995:2020))

colnames(mat.ssb) <- 0:17

rownames(mat.ssb) <- 1995:2020

for(y in 1:26){

tmp <- (as.data.frame(out.acu[[y]]$ssbByAge)[,c("age","Ton.mean")])

for(a in 1:18){

age <- a-1

if(!any(tmp$age[!is.na(tmp$age)] == age)) {

next

}

mat.ssb[y,a] <- tmp$`Ton.mean`[tmp$age == age & !is.na(tmp$age)]

}

}

## Figure 1: Number by age in time series ----

plot.cohort.year <- function(mat, age=1:17, survey.year=1995:2020, start.coh = 1995,
stop.coh=2016,col1=1,col2=1, addPlot=F){

## Define first cohort: start.co
```

```
## Does the matrix include year and age information? Add if not

mat <- mat[ ,colnames(mat) %in% age]

mat[mat == 1] <- NA

if(!is.null(rownames(mat))) survey.year <- as.numeric(rownames(mat))

mat <- mat[row.names(mat) %in% survey.year,]

age.mat <- matrix(age,nrow=nrow(mat),ncol=ncol(mat), byrow=T)

sur.mat <- matrix(survey.year,nrow=nrow(mat),ncol=ncol(mat), byrow=F)

coh.mat <- sur.mat - age.mat ####

if(addPlot == F) plot(sur.mat, mat, type="n", xlab="Survey year", ylab = "Log10 (abundance index)")

for(i.y in start.coh:stop.coh){

  lines(sur.mat[coh.mat== i.y], mat[coh.mat == i.y], col=i.y, lwd=1.5)

  text(sur.mat[coh.mat== i.y], mat[coh.mat == i.y], age.mat[coh.mat == i.y], col=i.y) ## M? skrive inn alder

}

## Plot cohort

fil1 <- paste0(path2,"Figure_1_Log10Number_by_age_in_time_series_CoastalCod.jpg")

jpeg(filename = fil1 , width = 18, height = 12, units = "cm",res=300)

plot.cohort.year(log10(mat.num), survey.year =1995:2020, age=1:17, start.coh=1995,stop.coh=2016)

dev.off()

# End

# Table 1: Abundance indices (mean bootstrap) ----

mat1.num <- as.data.frame(mat.num/(1e+6))

names(mat1.num) <- paste("Age", 0:(ncol(mat1.num)-1))

write.xlsx(mat1.num, file=paste0(path2,"Table_1_Abundance_indices_Coastal cod_Tot_millions.xlsx" ))

## Table 2: CV of abundance indices ----

mat.cv <- matrix(NA,ncol=18,nrow=length(1995:2020))

colnames(mat.cv) <- 0:17

rownames(mat.cv) <- 1995:2020

for(y in 1:26){
```

```
tmp <- (as.data.frame(out.acu[[y]]$tsnByAge)[,c("age","Ab.Sum.cv")])  
  
for(a in 1:18){  
  
  age <- a-1  
  
  if(!any(tmp$age[!is.na(tmp$age)] == age)) {  
  
    next  
  
  }  
  
  mat.cv[y,a] <- tmp$`Ab.Sum.cv`[tmp$age == age & !is.na(tmp$age)]  
  
}  
  
mat.cv1 <- as.data.frame(mat.cv)  
  
names(mat.cv1) <- paste("Age", 0:(ncol(mat.cv1)-1))  
  
write.xlsx(mat.cv1, file=paste0(path2,"Table_2_CV_Abundance_CoastalCod.xlsx" ))  
  
# Table 3: Biomass indices (mean bootstrap) ----  
  
mat1.tonn <- as.data.frame(mat.biom/(1e+3))  
  
names(mat1.tonn) <- paste("Age", 0:(ncol(mat1.tonn)-1))  
  
write.xlsx(mat1.tonn, file=paste0(path2,"Table_3_Biomass_indices_CoastalCod_kilotonnes.xlsx" ))  
  
#SSB indices  
  
mat1.tonn <- as.data.frame(mat.ssb/(1e+3))  
  
names(mat1.tonn) <- paste("Age", 0:(ncol(mat1.tonn)-1))  
  
write.xlsx(mat1.tonn, file=paste0(path2,"SSB_indices_CoastalCod_kilotonnes.xlsx" ))  
  
# Table 4: Mean length by age (cm)  
  
mat.len.age <- matrix(NA,ncol=18,nrow=length(1995:2020))  
  
colnames(mat.len.age) <- 0:17  
  
rownames(mat.len.age) <- 1995:2020  
  
for(y in 1:26){  
  
  tmp <- (as.data.frame(out.acu[[y]]$ind.length.by.age)[,c("age","ind.length.age.mean")])  
  
  for(a in 1:18){  
  
    age <- a-1  
  
    if(!any(tmp$age[!is.na(tmp$age)] == age)) {
```

```
next
}

mat.len.age[y,a] <- tmp$`ind.length.age.mean`[tmp$age == age & !is.na(tmp$age)]
}

}

mat.len.age1 <- as.data.frame(mat.len.age)

names(mat.len.age1) <- paste("Age", 0:(ncol(mat.len.age1)-1))

write.xlsx(mat.len.age1, file=paste0(path2,"Table_4_Length_By_Age_By_Survey_CoastalCod.xlsx" ))

# Table 5: Mean weight by age (g)

mat.weight.age <- matrix(NA, ncol=18, nrow=length(1995:2020))

colnames(mat.weight.age) <- 0:17

rownames(mat.weight.age) <- 1995:2020

for(y in 1:26){

tmp <- (as.data.frame(out.acu[[y]]$ind.weight.by.age)[,c("age","ind.weight.age.mean")])

for(a in 1:18){

age <- a-1

if(!any(tmp$age[!is.na(tmp$age)] == age)) {

next
}

mat.weight.age[y,a] <- tmp$`ind.weight.age.mean`[tmp$age == age & !is.na(tmp$age)]
}

}

mat.weight.age1 <- as.data.frame(mat.weight.age)

names(mat.weight.age1) <- paste("Age", 0:(ncol(mat.weight.age1)-1))

write.xlsx(mat.weight.age1, file=paste0(path2,"Table_5_Weight_By_Age_By_Survey_CoastalCod.xlsx" ))

#For assessment - maturity at age

mat.maturity.age <- matrix(NA, ncol=18, nrow=length(1995:2020))

colnames(mat.maturity.age) <- 0:17

rownames(mat.maturity.age) <- 1995:2020
```

```
for(y in 1:26){

tmp <- (as.data.frame(out.acu[[y]]$maturity.by.age)[,c("age","maturity.age.mean")])

for(a in 1:18){

age <- a-1

if(!any(tmp$age[!is.na(tmp$age)] == age)) {

next

}

mat.maturity.age[y,a] <- tmp`$maturity.age.mean`[tmp$age == age & !is.na(tmp$age)]

}

}

mat.maturity.age1 <- as.data.frame(mat.maturity.age)

names(mat.maturity.age1) <- paste("Age", 0:(ncol(mat.maturity.age1)-1))

write.xlsx(mat.maturity.age1, file=paste0(path2,"Maturity_By_Age_Acoustic_CoastalCod.xlsx" ))

#Table 6

mat.age2plus <- matrix(NA, ncol=3,nrow=length(1995:2020))

colnames(mat.age2plus) <- c( "Index_2plus" , "SD_2plus", "CV_2plus")

rownames(mat.age2plus) <- 1995:2020

for(y in 1:26){

mat.age2plus[y,1] <- (as.data.frame(out.acu[[y]]$tsnTotWithout01)[,c("Ab.Sum.mean")]/(1e+6))

mat.age2plus[y,2] <- (as.data.frame(out.acu[[y]]$tsnTotWithout01)[,c("Ab.Sum.sd")]/(1e+6))

mat.age2plus[y,3] <- (as.data.frame(out.acu[[y]]$tsnTotWithout01)[,c("Ab.Sum.cv")])

}

write.xlsx(mat.age2plus, file=paste0(path2,"Table_6_Index2+_w_sd-and-sv_CoastalCod.xlsx" ))

## End
```



## HAVFORSKNINGSINSTITUTTET

Postboks 1870 Nordnes

5817 Bergen

Tlf: 55 23 85 00

E-post: [post@hi.no](mailto:post@hi.no)

[www.hi.no](http://www.hi.no)